

A Mean Value Theorem for Closed Geodesics on Congruence Surfaces

Vladimir Lukianov
Tel-Aviv University

February 1, 2008

Abstract. We define a weighted multiplicity function for closed geodesics of given length on a finite area Riemann surface. These weighted multiplicities appear naturally in the Selberg trace formula, and in particular their mean square plays an important role in the study of statistics of the eigenvalues of the Laplacian on the surface.

In the case of the modular domain, E. Bogomolny, F. Leyvraz and C. Schmit gave a formula for the mean square, which was rigorously proved by M. Peter. In this paper we calculate the mean square of weighted multiplicities for some surfaces associated to congruence subgroups of the unit group of a rational quaternion algebra, in particular for congruence subgroups of the modular group. Remarkably, the result turns out to be a rational multiple of the mean square for the modular domain.

1 Introduction

The aim of the present work is to give a tool for finding the eigenvalue statistics of the hyperbolic Laplacian

$$\Delta = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$$

defined on a Riemann surface $\Omega = \Gamma \backslash \mathbb{H}$, for a discrete cofinite subgroup Γ of $PSL_2(\mathbb{R})$ of "congruence type" (see below).

Let $\lambda_0 = 0 < \lambda_1 \leq \lambda_2 \leq \dots$ be the discrete spectrum of the hyperbolic Laplacian on Ω . It is known [6] that it satisfies the Weyl's law:

$$N(T) := \#\{0 \leq r_j \leq T\} = \frac{\text{vol}(\Omega)}{4\pi} T^2 + O(T \ln T),$$

where $\lambda_j = \frac{1}{4} + r_j^2$. We define a smooth version of the counting function as follows. Let f be an even test function with the compactly supported smooth

Fourier transform $\widehat{f}$. Define

$$N_f(\tau) = \sum_{j \geq 0} f(L(r_j - \tau)) + f(L(-r_j - \tau)).$$

For example, if f is the characteristic function of the interval $(-1/2, 1/2)$, then $N_f(\tau)$ counts the number of r_j lying in the intervals $\pm(\tau - 1/2L, \tau + 1/2L)$.

Put

$$h(r) = f(L(r - \tau)) + f(L(-r - \tau))$$

then we can use the Selberg Trace Formula to express $N_f(\tau)$ as:

$$\begin{aligned} N_f(\tau) = \sum_{j \geq 0} h(r_j) = & \{\text{identity contribution}\} + \\ & + \{\text{hyperbolic contribution}\} + \\ & + \{\text{elliptic contribution}\} + \\ & + \{\text{parabolic and continuous spectrum contribution}\} \end{aligned} \quad (1)$$

We say that the element T of Γ is hyperbolic if $|trT| > 2$. Such an element is conjugate in $PSL_2(\mathbb{R})$ to a diagonal matrix $\begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix}$, where $\lambda > 1$ is real. For the element T , such that $|trT| = t$ we define the norm¹ of T to be

$$\mathcal{N}(T) = \lambda^2 = \left(\frac{t + \sqrt{t^2 - 4}}{2} \right)^2.$$

For us the most important term in formula (1) is the hyperbolic contribution term, which is defined explicitly as:

$$\sum_{t > 2\{T\}} \sum_{\substack{\text{hyperbolic} \\ |trT|=t}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{1/2} - \mathcal{N}(T)^{-1/2}} g(\ln \mathcal{N}(T)).$$

There are in general several conjugacy classes with $|trT| = t$. Define the weighted multiplicity function $\beta_\Gamma(t)$ by:

$$\beta_\Gamma(t) = \frac{1}{4} \sum_{\substack{\{T\} \subset \Gamma \\ T=T_0^k \text{ is hyperbolic} \\ |trT|=t}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{1/2} - \mathcal{N}(T)^{-1/2}},$$

where T_0 is a primitive hyperbolic element, that is T_0 is not a power of any other hyperbolic element. In this notation we can rewrite the hyperbolic contribution term as follows

$$\sum_{t \geq 2} \beta_\Gamma(t) g(\ln \mathcal{N}(T)),$$

¹The number λ^2 is called also a multiplier of T .

so the information about the weighted multiplicities will be useful for understanding the behavior of this term.

From the prime geodesic theorem it follows that the mean value of $\beta_\Gamma(t)$ is unity:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 < t \leq N} \beta_\Gamma(t) = 1.$$

Our main result is a computation of the mean square of the weighted multiplicities for the case that Γ is congruence subgroup of the unit group of a quaternion algebra over the rationals.

Let B be a quaternion algebra over $\mathbb{Q}$. Let

$$N_B : B \rightarrow \mathbb{Q}$$

$$Tr_B : B \rightarrow \mathbb{Q}$$

denote respectively the reduced norm and the reduced trace of the elements of B . If $\alpha \in B$, then

$$N_B(\alpha) = \alpha + \bar{\alpha}$$

$$Tr_B(\alpha) = \alpha \bar{\alpha}$$

where $\bar{\alpha}$ is the conjugate of α .

We recall that quaternion algebra over field F is either division algebra, or is isomorphic to $M_2(F)$.

Let d_B be the reduced discriminant of B , that is d_B is the product of all primes p , such that $B_p = B \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is a division algebra. Note that $d_B > 1$ if and only if B is a division algebra. We assume that B is indefinite, that is $B_\infty := B \otimes_{\mathbb{Q}} \mathbb{R} \cong M_2(\mathbb{R})$.

let R be an order in B , then for $\alpha \in R$ we have that both the reduced trace and the reduced norm of α are integers.

Fix now the isomorphism $B \otimes_{\mathbb{Q}} \mathbb{R} \cong M_2(\mathbb{R})$, and set

$$\Gamma_R = \{\alpha \in R \mid N_B(\alpha) = 1\}.$$

Under above isomorphism Γ_R is identified with a subgroup of $SL_2(\mathbb{R})$, which is a cofinite Fuchsian group, (and is co-compact if $d_B > 1$). Moreover, the traces of its elements are *integers*. In particular, one can consider $B = M_2(\mathbb{Q})$, and for the given natural Q the set

$$R = \left\{ \begin{bmatrix} a & b \\ Qc & d \end{bmatrix} \in M_2(\mathbb{Z}) \right\}.$$

Then B is indefinite quaternion algebra over $\mathbb{Q}$ and R is an order in B . Here

$$\Gamma_R = \Gamma_0(Q) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z}) \mid c \equiv 0 \pmod{Q} \right\}. \quad (2)$$

Remark 1.1 *We will often identify Γ_R with its image in the projective group $PSL_2(\mathbb{R})$.*

We are now able to state our main theorems:

Theorem 1.2 *Let*

$$\beta_Q(t) = \frac{1}{4} \sum_{\substack{\{T\} \subset \Gamma_0(Q) \\ T=T_0^k \text{ is hyperbolic} \\ |trT|=t}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{1/2} - \mathcal{N}(T)^{-1/2}},$$

where $\Gamma_0(Q)$ is the congruence subgroup of $SL_2(\mathbb{Z})$, defined in (2). Then for the squarefree odd Q

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 < t \leq N} \beta_Q^2(t) = C_1 \prod_{q|Q} \frac{2(q^2 - q - 1)(q + 1)^2}{q(q^3 + q^2 - q - 3)} = \kappa_Q.$$

Here

$$C_1 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 < t \leq N} \beta_1^2(t) = \frac{1015}{864} \prod_{p \neq 2} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p + 1)} = 1.328...$$

which is the mean square of the weighted multiplicities for the modular domain ($Q = 1$), proved by M. Peter [10], following a conjecture of Bogomolny et al [3].

Theorem 1.3 *Let B be an indefinite division quaternion algebra over $\mathbb{Q}$ with the maximal order R , and discriminant d_B . Then for*

$$\beta_R(t) = \frac{1}{4} \sum_{\substack{\{T\} \subset \Gamma_R \\ T=T_0^k \text{ is hyperbolic} \\ |trT|=t}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{1/2} - \mathcal{N}(T)^{-1/2}}$$

the mean square of weighted multiplicities is

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n) = C_1 \cdot \prod_{p|d_B} \frac{2(p^3 - 1)(p - 1)}{p(p^3 + p^2 - p - 3)}.$$

We now explain the method of proof. As a first step, we express the weighted multiplicities in terms of Dirichlet's L -functions. For $\Gamma_0(Q)$ the expression is

$$\beta_Q(t) = \sum_{\substack{D, v \geq 1 \\ D \text{ is a discriminant} \\ Dv^2 = t^2 - 4}} \frac{1}{v} L(1, \chi_D) \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right) \begin{matrix} q^2 \mid D \\ q^2 \nmid D \end{matrix} \right\}. \quad (3)$$

This is proved in section 3 by connecting the weighted multiplicities with class numbers and using Dirichlet's class number formula.

The principal tool in our approach, following Peter, is that the formula (3) displays the weighted multiplicity $\beta_Q(t)$ as a "limit periodic function" in a suitable sense (see section 5 for background on these). To show this, we approximate the L -functions by a finite Euler product using a zero-density theorem, in a certain semi-norm coming from the theory of limit periodic functions (section 6). For computing mean squares, this suffices and allows us to use Parseval's equality in this setting to express the mean square as

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq t \leq N} \beta_Q^2(t) &= \sum_{b \geq 1} \sum_{\substack{1 \leq a \leq b \\ \gcd(a,b)=1}} \left| \widehat{\beta_Q} \left(\frac{a}{b} \right) \right|^2 = \\ &= \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} \sum_{\substack{1 \leq a \leq p^c \\ a \not\equiv 0 \pmod{p}}} \left| \widehat{\beta_{(p,Q)}} \left(\frac{a}{p^c} \right) \right|^2 \right), \end{aligned}$$

where $\widehat{\beta}$ are the Fourier coefficients of β , defined in section 5.

We then carry out a lengthy calculation of the Fourier coefficients in section 7.2, finally ending up with rather complicated expressions described in Theorem 7.2.

The result is that the mean square is an Euler product

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n) = \prod_{p \text{ - prime}} M_p(Q),$$

with

$$M_p(Q) = 1 + \sum_{c \geq 1} A_Q(p^c),$$

and $A_Q(p^c)$ is given by (12), section 7.3. We evaluate the sum $M_p(Q)$ over prime powers as a rational function of p and find that it depends on divisibility of Q by p , in particular, for $p \nmid Q$, $p \neq 2$,

$$M_p(Q) = M_p(1) = \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p + 1)}.$$

This will prove Theorem 1.2.

In section 8 we sketch this procedure for the case of the unit group of the maximal order of a quaternion algebra (theorem 1.3).

An application:

The computation of the mean square is used to study the statistics of $N_f(\tau)$: by Weyl's law we expect $N_f(\tau)$ to be asymptotically equal to a multiple of τ/L . To

study the fluctuations around this expectation one may consider τ as a random variable.

Then define an averaging operator:

$$\langle F \rangle_T = \frac{1}{T} \int_T^{2T} F(\tau) d\tau$$

Studying the moments of N_f shows that if $L \rightarrow \infty$, but $L = o(\ln T)$, then its limiting value distribution is Gaussian with the mean

$$\langle N_f(\tau) \rangle_T = \frac{\tau}{L} \frac{2 \text{vol}(\Omega)}{4\pi} \int_{-\infty}^{\infty} f(x) dx$$

and the variance

$$\left\langle (N_f(\tau) - \langle N_f(\tau) \rangle_T)^2 \right\rangle_T = \frac{4\kappa}{\pi L} \int_0^{\infty} \hat{f}^2(u) e^{\pi L u} du,$$

where κ is the mean square of the weighted multiplicities. See [14], [9]. Thus our computation of κ for Γ of congruence type yields the value distribution of N_f .

Acknowledgement. This work was supported in part by Gutwirth Scholarship and by the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities. It was carried out as part of the author's PHD thesis at Tel Aviv University, under the supervision of prof. Ze'ev Rudnick.

2 Preliminaries on orders in quaternion algebras

First of all we recall that a ring B with unity is called an algebra of dimension n over a field F , if the following three conditions are satisfied:

- 1° $B \supset F$, and $1_B = 1_F$;
- 2° any element of F commutes with all elements of B ;
- 3° B is a vector space over F of dimension n .

Definition 2.1 *Let B be a finite dimensional algebra over $\mathbb{Q}$ with identity element 1_B . A subset $R \subset B$ is called an order in B if the following conditions are satisfied:*

- 1° R is a finitely generated $\mathbb{Z}$ -module;
- 2° R contains basis of B over $\mathbb{Q}$;
- 3° R is a subring of B and $1_B \in R$.

For any finite dimensional algebra B over $\mathbb{Q}$ and for each place v of $\mathbb{Q}$ (i.e. $v = \infty$ or $v = p$ a prime) we define $B_v := B \otimes_{\mathbb{Q}} \mathbb{Q}_v$. B_v is an algebra over $\mathbb{Q}_v$ of dimension $\dim_{\mathbb{Q}} B$. For $v = p$ a prime, orders in B_p defined as in the previous definition, replacing $\mathbb{Q}$ and $\mathbb{Z}$ by $\mathbb{Q}_p$ and $\mathbb{Z}_p$. If R is an order of B , we get an order in B_p by the definition

$$R_p := R \otimes_{\mathbb{Z}} \mathbb{Z}_p = [\text{the closure of } R \text{ in } B_p].$$

We also recall the definition of adele ring B_A [13, pp. 197-198] and [17, p. 62]. As a set this is

$$B_A = \{(b_v) \in \prod_v B_v \mid b_p \in R_p \text{ for almost all primes } p\},$$

where R is any order in B .

Definition 2.2 *A nonzero integer d is called a fundamental discriminant if $d \equiv 1 \pmod{4}$, d square free, $d \neq 1$; or $d \equiv 0 \pmod{4}$, $\frac{d}{4} \not\equiv 1 \pmod{4}$, $\frac{d}{4}$ squarefree.*

Consider now a quadratic extension of $\mathbb{Q}$ arising as the splitting field over $\mathbb{Q}$ of a polynomial $P(X) = X^2 - tX + 1$, $t \in \mathbb{Z}$. Notice that $P(X)$ is irreducible iff $\sqrt{t^2 - 4} \notin \mathbb{Q}$. In this case we can write $t^2 - 4 = l^2 d$, $l \in \mathbb{Z}^+$, d is a fundamental discriminant. The splitting field of $P(X)$ is $\mathbb{Q}(\sqrt{d})$ and the zeros of $P(X)$ are

$$x_1 = \frac{t}{2} + l \frac{\sqrt{d}}{2}, \quad x_2 = \frac{t}{2} - l \frac{\sqrt{d}}{2}.$$

For each fixed fundamental discriminant d we let for each $f \in \mathbb{Z}^+$

$$\mathfrak{r}[f] := \mathbb{Z} + f\omega\mathbb{Z}, \quad \text{where } \omega = \frac{d + \sqrt{d}}{2}.$$

Then orders in $\mathbb{Q}(\sqrt{d})$ are precisely $\mathfrak{r}[f]$, $f = 1, 2, 3, \dots$ [2, pp. 48-49], and in particular $\mathfrak{r}[1]$ is the unique maximal order. [5, pp. 146-147]. The index $[\mathfrak{r}[1] : \mathfrak{r}[f]] = f$, and we call $\mathfrak{r}[f]$ the order of index f in $\mathbb{Q}(\sqrt{d})$. Note that $\mathfrak{r}[f_1] \subset \mathfrak{r}[f_2]$ iff $f_2 \mid f_1$.

For $P(X)$ as above, the roots x_1, x_2 of $P(X)$ satisfy

$$\mathbb{Z} + x_1\mathbb{Z} = \mathbb{Z} + x_2\mathbb{Z} = \mathfrak{r}[l] \quad (t^2 - 4 = l^2 d).$$

Let $K = \mathbb{Q}(\sqrt{d})$, d as above, then

$$K_{\infty} = \mathbb{Q}(\sqrt{d}) \otimes_{\mathbb{Q}} \mathbb{R} \simeq \left\{ \begin{array}{ll} \mathbb{R} \oplus \mathbb{R} & , d > 0 \\ \mathbb{C} & , d < 0 \end{array} \right\},$$

$$K_p = \mathbb{Q}(\sqrt{d}) \otimes_{\mathbb{Q}} \mathbb{Q}_p \simeq \begin{cases} \mathbb{Q}_p \oplus \mathbb{Q}_p & , \left(\frac{d}{p}\right) = 1 \\ \mathbb{Q}_p(\sqrt{d}) & , \left(\frac{d}{p}\right) = -1 \end{cases}.$$

The distinct orders in K_p are

$$\mathfrak{o}[p^n] := \mathbb{Z}_p + p^n \omega \mathbb{Z}_p, n = 0, 1, 2, \dots \text{ and } \omega = \frac{d + \sqrt{d}}{2} \text{ as above.}$$

We have

$$\mathfrak{o}[1] \supset \mathfrak{o}[p] \supset \mathfrak{o}[p^2] \supset \dots \text{ and } [\mathfrak{o}[1] : \mathfrak{o}[p^n]] = p^n.$$

Combining the definitions of order in K and in K_p we get

$$(\mathfrak{r}[f])_p = \mathfrak{o}[p^n], \text{ where } n = \text{ord}_p f.$$

For any order $\mathfrak{r}$ in $K = \mathbb{Q}(\sqrt{d})$ we define

$$\begin{aligned} \mathfrak{r}_\infty &:= K_\infty \\ \mathfrak{r}_A &:= \prod_v \mathfrak{r}_v \subset K_A \\ \mathfrak{r}_{\infty+}^\times &:= \{x \in K_\infty^\times \mid N(x) > 0\} \\ \mathfrak{r}_{A+}^\times &:= \{(\alpha_v) \in \mathfrak{r}_A^\times \mid N(\alpha_\infty) > 0\} = \mathfrak{r}_{\infty+}^\times \times \prod_p \mathfrak{r}_p^\times \subset \mathfrak{r}_A^\times \end{aligned}$$

Here N denotes the extension of the norm $N : K \longrightarrow \mathbb{Q}$ to the regular norm in the algebra K_∞ over $\mathbb{Q}_\infty = \mathbb{R}$. [17, p.53].

The subgroup $\mathfrak{r}_{A+}^\times \cdot K^\times$ has finite index in $K_A^\times$, and we set

$$h(\mathfrak{r}) := [K_A^\times : (\mathfrak{r}_{A+}^\times \cdot K^\times)],$$

the class number of $\mathfrak{r}$.

For $K = \mathbb{Q}(\sqrt{d})$ as above one can also show [1, Chapter 5.2] that $h(\mathfrak{r}[f])$ is equal to the number of inequivalent primitive (and if $d < 0$, positive) quadratic forms $ax^2 + bxy + cy^2$ with discriminant $b^2 - 4ac = df^2$. In other words $h(\mathfrak{r}[f]) = h(df^2)$, where the right-hand side is as in [8, vol I, pp. 127-].

3 Expression of weighted multiplicities function in the terms of Dirichlet's L -functions

In this section we define the weighted multiplicities function $\beta_Q(t)$ and express it in the terms of Dirichlet's L -functions. It is a necessary step, which allows to analyze the behavior of $\beta_Q(t)$. The result is stated in Theorem 3.1, proof of which takes the rest of the section.

Let $T \in SL_2(\mathbb{Z})$, such that $|t| = |tr(T)| > 2$. For such T the equation $\det(T - \lambda I) = 0$ has two different solutions: $\lambda_{1,2} = \frac{t \pm \sqrt{t^2 - 4}}{2}$, and so T is diagonalizable in $\mathbb{R}$. Such T is called hyperbolic. Define $\mathcal{N}(T) := \lambda^2$, where $|\lambda| > 1$. Hence $\mathcal{N}(T) = \frac{(|t| + \sqrt{t^2 - 4})^2}{4}$.

Let Q be an odd squarefree integer, and let

$$\Gamma_0(Q) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z}) \mid c \equiv 0 \pmod{Q} \right\}$$

be the congruence subgroup of $SL_2(\mathbb{Z})$. Define the weighted multiplicities function

$$\beta_Q(t) := \frac{1}{4} \sum_{\substack{\{T\} \in \Gamma_0(Q) \\ |tr(T)|=t \\ \text{hyperbolic, doesn't} \\ \text{fix cusp}}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{\frac{1}{2}} - \mathcal{N}(T)^{-\frac{1}{2}}},$$

where the sum is taken over conjugacy classes of $\Gamma_0(Q)$.

Theorem 3.1 For $2 < |t| \in \mathbb{Z}$

$$\beta_Q(t) = \sum_{\substack{D, v \geq 1 \\ D \text{ is a discriminant} \\ Dv^2 = t^2 - 4}} \frac{1}{v} L(1, \chi_D) \cdot \prod_{q|Q} \left\{ 1 + \begin{matrix} 2, & q^2 \mid D \\ \left(\frac{D}{q}\right) & q^2 \nmid D \end{matrix} \right\},$$

and $\beta_Q(t) = 0$, otherwise. Here $(\cdot)$ is a Legendre's symbol and χ_D is a quadratic character.

We start from noting that

$$\Gamma_0(Q) = \bigsqcup_{t \in \mathbb{Z}} H_t,$$

where

$$H_t = \{T \in \Gamma_0(Q) \mid tr(T) = t\}.$$

So we can write (the change from 1/4 to 1/2 comes from not collecting together t and $-t$)

$$\beta_Q(t) = \frac{1}{2} \sum_{\substack{T \in H_t / \Gamma_0(Q) \\ \text{hyperbolic, doesn't} \\ \text{fix cusp}}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{\frac{1}{2}} - \mathcal{N}(T)^{-\frac{1}{2}}} \quad (4)$$

Here $H_t / \Gamma_0(Q)$ is the set of $\Gamma_0(Q)$ -conjugacy classes of H_t .

Lemma 3.2 *For a hyperbolic element $T \in H_t$ (i.e. $t^2 - 4 > 0$) no fixed point of T is a cusp of $\Gamma_0(Q)$.*

Proof. Note, that the cusps of $\Gamma_0(Q)$ are exactly the points in $\{i\infty\} \cup \mathbb{Q}$, by [13, cor. 1.5.5, th. 4.1.3(2)]. Assume $T = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. First, $c \neq 0$, since we will get T parabolic. So, the fixpoints of T are the two real solutions to $\frac{ax+b}{cx+d} = x \Leftrightarrow x = \frac{a-d}{2c} \pm \frac{\sqrt{t^2-4}}{2|c|}$, which are both irrational, since $\sqrt{t^2-4} \notin \mathbb{Q}$. ■

Now we want to know when in the sum (4), H_t is not the empty set.

Lemma 3.3 *Let $t \in \mathbb{Z}$, such that $\sqrt{t^2-4} \notin \mathbb{Q}$, and write $t^2 - 4 = l^2 d$, $l \in \mathbb{Z}^+$, d is a fundamental discriminant. Then H_t is not empty iff for all primes q divide Q we have $q \mid l$ or $\left(\frac{d}{q}\right) \neq -1$.*

Proof. $H_t \neq \emptyset$ iff there some $a, b, c \in \mathbb{Z}$, such that $\det \begin{bmatrix} a & b \\ Qc & t-a \end{bmatrix} = 1$. Hence $H_t \neq \emptyset$ iff there is some $a \in \mathbb{Z}$, such that $a(t-a) \equiv 1 \pmod{Q} \Leftrightarrow (2a-t)^2 \equiv t^2-4 \pmod{Q} \Leftrightarrow (2a-t)^2 \equiv l^2 d \pmod{Q}$. Since Q is odd squarefree, the last congruence is solvable iff $(2a-t)^2 \equiv l^2 d \pmod{q}$ for all primes $q \mid Q$. Or equivalently: the last congruence is solvable iff $q \mid l$ or $\left(\frac{d}{q}\right) \neq -1$, for all $q \mid Q$. ■

Note, that for $T \in H_t$

$$\mathcal{N}(T)^{\frac{1}{2}} - \mathcal{N}(T)^{-\frac{1}{2}} = \frac{|t| + \sqrt{t^2-4}}{2} - \frac{2}{|t| + \sqrt{t^2-4}} = \sqrt{t^2-4}.$$

So, for fixed $t \in \mathbb{Z}$, such that $\sqrt{t^2-4} \notin \mathbb{Q}$, $t^2 - 4 > 0$, $H_t \neq \emptyset$ we can write (4) in the form

$$\beta_Q(t) = \frac{1}{2} \sum_{T \in H_t / \Gamma_0(Q)} \frac{\ln \mathcal{N}(T_0)}{\sqrt{t^2-4}} \quad (5)$$

We will enumerate the $\Gamma_0(Q)$ -conjugacy classes in H_t , and for each conjugacy class will get the number $\mathcal{N}(T_0)$.

Fix some $T_t \in H_t$, and take the polynomial $P(X) = X^2 - tX + 1$, i.e. the characteristic polynomial of T_t , and so $P(T_t) = 0$. $P(X)$ is irreducible, since $\sqrt{t^2-4} \notin \mathbb{Q}$. Hence $\mathbb{Q}[T_t] \cong \mathbb{Q}(\sqrt{d})$, where $t^2 - 4 = l^2 d$.

Define the set $C(T_t) := \{\delta T_t \delta^{-1} \mid \delta \in GL_2(\mathbb{Q})\}$. Then $H_t = R \cap C(T_t)$, where $R = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{Z}) \mid Q \mid c \right\}$ is an order in $M_2(\mathbb{Q})$.

For any $\delta \in GL_2(\mathbb{Q})$, $(\mathbb{Q}[T_t] \cap \delta R \delta^{-1})$ is an order in $\mathbb{Q}[T_t]$, by the fact, that if A' is a subalgebra of A , and $\mathcal{O}$ is an order in A , then $A' \cap \mathcal{O}$ is an order in A' . For any order $\mathfrak{r}$ in $\mathbb{Q}[T_t]$ define $C(T_t, \mathfrak{r}) := \{\delta T_t \delta^{-1} \mid \delta \in GL_2(\mathbb{Q}), \mathbb{Q}[T_t] \cap \delta^{-1} R \delta = \mathfrak{r}\}$. Following [16], [13] we have a

Lemma 3.4

$$C(T_t) = \bigsqcup_{f=1}^{\infty} C(T_t, \mathfrak{r}[f]),$$

$$H_t = \bigsqcup_{f \mid l} C(T_t, \mathfrak{r}[f]),$$

and each $C(T_t, \mathfrak{r}[f])$ is closed under $\Gamma_0(Q)$ -conjugation.²

Proof. Clearly, $C(T_t) = \bigcup_{f=1}^{\infty} C(T_t, \mathfrak{r}[f])$, since there are no other orders than $\mathfrak{r}[1], \mathfrak{r}[2], \dots$ in $\mathbb{Q}[T_t]$. To prove disjointedness, we must show that if $\delta_1 T_t \delta_1^{-1} = \delta_2 T_t \delta_2^{-1}$ ($\delta_1, \delta_2 \in GL_2(\mathbb{Q})$), then $\delta_1^{-1} R \delta_1$, and $\delta_2^{-1} R \delta_2$ have the same intersection with $\mathbb{Q}[T_t]$. From $\delta_1 T_t \delta_1^{-1} = \delta_2 T_t \delta_2^{-1}$ we get $\delta_1^{-1} \delta_2 \in \mathbb{Q}[T_t]$, since if an element of $M_2(\mathbb{Q})$ commutes with T_t , it is in $\mathbb{Q}[T_t]$, by [13, Lemma 5.2.2(3)], and thus

$$\mathbb{Q}[T_t] \cap \delta_2^{-1} R \delta_2 = (\delta_1^{-1} \delta_2) (\mathbb{Q}[T_t] \cap \delta_2^{-1} R \delta_2) (\delta_1^{-1} \delta_2)^{-1} = \mathbb{Q}[T_t] \cap \delta_1^{-1} R \delta_1.$$

This prove the first relation.

Next, for any order $\mathfrak{r}$ in $\mathbb{Q}[T_t]$ and any $\delta \in GL_2(\mathbb{Q})$, such that $\delta T_t \delta^{-1} \in C(T_t, \mathfrak{r})$, we have :

$$T_t \in \mathfrak{r} \Leftrightarrow T_t \in \mathbb{Q}[T_t] \cap \delta^{-1} R \delta \Leftrightarrow T_t \in \delta^{-1} R \delta \Leftrightarrow \delta T_t \delta^{-1} \in R \Leftrightarrow \delta T_t \delta^{-1} \in H_t.$$

In other words :

$$\begin{aligned} &\text{if } T_t \in \mathfrak{r}, \text{ then } C(T_t, \mathfrak{r}) \subset H_t; \\ &\text{if } T_t \notin \mathfrak{r}, \text{ then } C(T_t, \mathfrak{r}) \cap H_t = \emptyset. \end{aligned}$$

Note that $T_t \in \mathfrak{r}[l]$, where $t^2 - 4 = l^2 d$, $l \in \mathbb{Z}^+$. So the orders of $\mathfrak{r}$ in $\mathbb{Q}[T_t]$ that contain T_t are exactly $\mathfrak{r}[f]$, such that $\mathfrak{r}[f] \supset \mathfrak{r}[l]$, that is, such that $f \mid l$. By definition $H_t \subset C(T_t)$, and so from the first relation the second relation follows.

² $\mathfrak{r}[f] := \mathbb{Z} + f w \mathbb{Z}$, is an all orders in $\mathbb{Q}(\sqrt{d})$, where $f \in \mathbb{Z}^+$, $w = \frac{d + \sqrt{d}}{2}$. Since $\mathbb{Q}(\sqrt{d}) \cong \mathbb{Q}[T_t]$ we use the symbol $\mathfrak{r}[f]$ for the corresponding order in $\mathbb{Q}[T_t]$.

Finally, $\gamma^{-1}R\gamma = R$, for any $\gamma \in R^\times$. So we get that each $C(T_t, \mathfrak{r})$ is closed under $R^\times$ -conjugation, and in particular under $\Gamma_0(Q)$ -conjugation. ■

Let $\varepsilon_d = \frac{x + y\sqrt{d}}{2}$ be the proper fundamental unit in $\mathbb{Q}(\sqrt{d})$ ((x, y) is the positive integer solution to $x^2 - dy^2 = 4$, for which $y > 0$ is minimal). Define

$$\mathfrak{r}[f]^1 := \{\alpha \in \mathfrak{r}[f] \mid N(\alpha) = 1\},$$

the units of the order $\mathfrak{r}[f]$. Since

$$\mathfrak{r}[1]^1 = \{\pm \varepsilon_d^k \mid k \in \mathbb{Z}\},$$

we have

$$\mathfrak{r}[f]^1 = \left\{ \pm \left(\varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]} \right)^k \mid k \in \mathbb{Z} \right\}.$$

Lemma 3.5 *If T is hyperbolic, i.e. $d > 0$, then $\mathcal{N}(T_0) = \varepsilon_d^{2[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}$.*

Proof. Choose some $\delta \in GL_2(\mathbb{Q})$, such that $T = \delta T_t \delta^{-1}$, and $\mathbb{Q}[T_t] \cap \delta R \delta^{-1} = \mathfrak{r}$. Then $Z_{\Gamma_0(Q)}(T)$, the centralizer of T in $\Gamma_0(Q)$ is

$$Z_{\Gamma_0(Q)}(T) = \mathbb{Q}[T] \cap \Gamma_0(Q) = \delta(\mathbb{Q}[T_t] \cap \delta^{-1}\Gamma_0(Q)\delta)\delta^{-1} = \delta \mathfrak{r}[f] \delta^{-1}$$

by [13, lemma 5.2.2(3)]. Since

$$\mathfrak{r}[f]^1 = \left\{ \pm \left(\varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]} \right)^k \mid k \in \mathbb{Z} \right\},$$

for $d > 0$ we can take for T_0 the image of $\varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}$ under the composite of the two isomorphisms $\mathbb{Q}[T_t] \cong \mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}[T_t] \ni S \mapsto \delta S \delta^{-1} \in \mathbb{Q}[T]$. Then

$$tr_{M_2(\mathbb{Q})}(T_0) = tr \left(\varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]} \right).$$

If we will denote $\varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]} = \alpha + \beta\sqrt{d} \in \mathbb{Q}(\sqrt{d})$, note, that $\alpha, \beta \geq 0$, and $N(\alpha + \beta\sqrt{d}) = \alpha^2 - d\beta^2 = 1$. We have $tr(\alpha + \beta\sqrt{d}) = t$, and so

$$\begin{aligned} \mathcal{N}(T_0) &= \frac{(|t| + \sqrt{t^2 - 4})^2}{4} = \frac{(2|\alpha| + \sqrt{4\alpha^2 - 4})^2}{4} = \\ &= \frac{(2|\alpha| + \sqrt{4(1 + d\beta^2) - 4})^2}{4} = \frac{(2|\alpha| + 2|\beta|\sqrt{d})^2}{4} = \\ &= (|\alpha| + |\beta|\sqrt{d})^2 = (\alpha + \beta\sqrt{d})^2 = \varepsilon_d^{2[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}. \end{aligned}$$

■

Now we can rewrite (5) in the form

$$\beta_Q(t) = \sum_{f|l} |C(T_t, \mathfrak{r}[f]) / \Gamma_0(Q)| \frac{\ln \varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}}{\sqrt{t^2 - 4}},$$

where we write $t^2 - 4 = dl^2$ with d a fundamental discriminant and $l \geq 1$.

For the quantities $|C(T_t, \mathfrak{r}[f]) / \Gamma_0(Q)|$ we refer to the [13, §6.6], [16]:

Proposition 3.6 *Let Q be squarefree and $h(df^2)$ the narrow class number of $\mathbb{Q}(\sqrt{d})$. Then*

$$|C(T_t, \mathfrak{r}[f]) / \Gamma_0(Q)| = h(df^2) \cdot \begin{cases} 2, & \text{if } d < 0 \\ 1, & \text{if } d > 0 \end{cases} \cdot \prod_{p|Q} \begin{cases} 2, & p \mid f \\ 1 + \left(\frac{d}{p}\right), & p \nmid f \end{cases}.$$

Corollary 3.7

$$\beta_Q(t) = \sum_{f|l} h(df^2) \frac{\ln \varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}}{\sqrt{t^2 - 4}} \cdot \prod_{q|Q} \begin{cases} 2, & q \mid f \\ 1 + \left(\frac{d}{q}\right), & q \nmid f \end{cases},$$

where we write $t^2 - 4 = dl^2$ with d a fundamental discriminant and $l \geq 1$.

Proof. Immediately from proposition 3.6. ■

We can continue the process

$$\begin{aligned} \beta_Q(t) &= \sum_{f|l} h(df^2) \frac{\ln \varepsilon_d^{[\mathfrak{r}[1]^1 : \mathfrak{r}[f]^1]}}{\sqrt{t^2 - 4}} \cdot \prod_{q|Q} \begin{cases} 2, & q \mid f \\ 1 + \left(\frac{d}{q}\right), & q \nmid f \end{cases} = \\ &= \sum_{f|l} \frac{h(df^2) \ln \varepsilon_{df^2}}{l\sqrt{d}} \cdot \prod_{q|Q} \begin{cases} 2, & q \mid f \\ 1 + \left(\frac{d}{q}\right), & q \nmid f \end{cases} = \\ &= \sum_{\substack{f|l \\ (D=df^2)}} \frac{h(D) \ln \varepsilon_D}{l \cdot \frac{\sqrt{D}}{f}} \cdot \prod_{q|Q} \begin{cases} 2, & q \mid f \\ 1 + \left(\frac{d}{q}\right), & q \nmid f \end{cases}. \end{aligned}$$

Now using Dirichlet's class number formula $h(D) \ln \varepsilon_D = \sqrt{D} L(1, \chi_D)$, we get

$$\begin{aligned}
\beta_Q(t) &= \sum_{\substack{f|l \\ (D=df^2)}} L(1, \chi_D) \frac{f}{l} \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q|f \\ q \nmid f \end{matrix} \right\} = \\
&= \sum_{\substack{D, v \geq 1 \\ Dv^2 = t^2 - 4}} \frac{1}{v} L(1, \chi_D) \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q|f \\ q \nmid f \end{matrix} \right\} = \\
&= \sum_{\substack{D, v \geq 1 \\ Dv^2 = t^2 - 4}} \frac{1}{v} L(1, \chi_D) \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 | D \\ q^2 \nmid D \end{matrix} \right\},
\end{aligned}$$

and the theorem 3.1 follows.

Remark 3.8 Here D is a discriminant, i.e. $D \equiv 0, 1 \pmod{4}$. We assume it from now on.

4 Factorization of weighted multiplicities

Here we factorize (lemma 4.1) the weighted multiplicities function as a finite products of a local terms, defined below.

We have

$$\beta_Q(n) = \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4}} \frac{1}{v} L(1, \chi_D) \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 | D \\ q^2 \nmid D \end{matrix} \right\}.$$

We use now the Euler product formula for $L(1, \chi_D)$. For $n \geq 3, P \geq Q$, define

$$\beta_{P,Q}(n) := \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ p|v \implies p \leq P}} \left(\frac{1}{v} \prod_{p \leq P} \left(1 - \frac{\chi_D(p)}{p} \right)^{-1} \right) \cdot \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 | D \\ q^2 \nmid D \end{matrix} \right\}.$$

Note, that

$$\begin{aligned}
q^2 | D &\iff q^2 | (n^2 - 4)v^{-2} \iff q^2 | (n^2 - 4) \prod_{\substack{p \leq P \\ p|v}} p^{-2b_p} \\
&\iff \left\{ \begin{matrix} q^2 | n^2 - 4, & b_q = 0 \\ q^2 | (n^2 - 4)q^{-2b_q}, & b_q \neq 0 \end{matrix} \right\} \iff q^2 | (n^2 - 4)q^{-2\text{ord}_q v}.
\end{aligned}$$

Hence

$$\beta_{P,Q}(n) = \sum_{\substack{D,v \geq 1 \\ Dv^2 = n^2 - 4 \\ p|v \implies p \leq P}} \left(\frac{1}{v} \prod_{p \leq P} \left(1 - \frac{\chi_D(p)}{p} \right)^{-1} \right) \times \\ \times \prod_{q|Q} \left\{ 1 + \left(\frac{2}{(n^2 - 4)q^{-2\text{ord}_q v}} \right) \begin{array}{l} , \quad q^2 \mid (n^2 - 4)q^{-2\text{ord}_q v} \\ , \quad q^2 \nmid (n^2 - 4)q^{-2\text{ord}_q v} \end{array} \right\}.$$

Define a function

$$\beta_{(p,Q)}(n) := \sum_{b \geq 0} \frac{1}{p^b} \left(1 - \frac{1}{p} \chi_{(n^2 - 4)p^{-2b}}(p) \right)^{-1} \cdot \mathbb{I}_{p^b}(n),$$

where

$$\mathbb{I}_{p^b}(n) := \left\{ \begin{array}{ll} 1 & , n^2 = 4 \pmod{p^{2b}} \\ 0 & , \text{else} \end{array} \right\}, \text{ for } p \neq 2, p \nmid Q$$

$$\mathbb{I}_{2^b}(n) := \left\{ \begin{array}{ll} 1 & , n^2 = 4 \pmod{2^{2b}} \quad , (n^2 - 4)2^{-2b} \text{ is a discriminant} \\ 0 & , \text{else} \end{array} \right\},$$

and for $q \mid Q$

$$\mathbb{I}_{q^b}(n) := \left\{ \begin{array}{lll} 2 & , n^2 = 4 \pmod{q^{2b}} & , q^2 \mid (n^2 - 4)q^{-2b} \\ 1 + \left(\frac{(n^2 - 4)q^{-2b}}{q} \right) & , n^2 = 4 \pmod{q^{2b}} & , q^2 \nmid (n^2 - 4)q^{-2b} \\ 0 & , \text{else} \end{array} \right\}.$$

Lemma 4.1 For Q odd squarefree and $P \geq Q$ we have

$$\beta_{P,Q}(n) = \prod_{\substack{p \leq P \\ p \nmid Q}} \beta_{(p,Q)}(n) \cdot \prod_{q|Q} \beta_{(q,Q)}(n).$$

Proof.

$$\prod_{\substack{p \leq P \\ p \nmid Q}} \beta_{(p,Q)}(n) \cdot \prod_{q|Q} \beta_{(q,Q)}(n) = \prod_{\substack{p \leq P \\ p \nmid Q}} \left(\sum_{b \geq 0} \frac{1}{p^b} \left(1 - \frac{1}{p} \chi_{(n^2 - 4)p^{-2b}}(p) \right)^{-1} \cdot \mathbb{I}_{p^b}(n) \right) \times \\ \prod_{q|Q} \left(\sum_{b \geq 0} \frac{1}{q^b} \left(1 - \frac{1}{q} \chi_{(n^2 - 4)q^{-2b}}(q) \right)^{-1} \cdot \mathbb{I}_{q^b}(n) \right).$$

After opening the brackets, we will get the sum of terms of the form:

$$\begin{aligned}
& \frac{1}{p_1^{b_1}} \left(1 - \frac{1}{p_1} \chi_{(n^2-4)p_1^{-2b_1}}(p_1) \right)^{-1} \mathbb{I}_{p_1^{b_1}}(n) \cdot \frac{1}{p_2^{b_2}} \left(1 - \frac{1}{p_2} \chi_{(n^2-4)p_2^{-2b_2}}(p_2) \right)^{-1} \mathbb{I}_{p_2^{b_2}}(n) \cdot \dots \cdot \\
& \frac{1}{p_k^{b_k}} \left(1 - \frac{1}{p_k} \chi_{(n^2-4)p_k^{-2b_k}}(p_k) \right)^{-1} \mathbb{I}_{p_k^{b_k}}(n) \cdot \frac{1}{q_1^{b_{q_1}}} \left(1 - \frac{1}{q_1} \chi_{(n^2-4)q_1^{-2b_{q_1}}}(q_1) \right)^{-1} \mathbb{I}_{q_1^{b_{q_1}}}(n) \cdot \dots \cdot \\
& \frac{1}{q_l^{b_{q_l}}} \left(1 - \frac{1}{q_l} \chi_{(n^2-4)q_l^{-2b_{q_l}}}(q_l) \right)^{-1} \mathbb{I}_{q_l^{b_{q_l}}}(n).
\end{aligned}$$

Therefore, since $P \geq Q$ we have

$$\begin{aligned}
& \prod_{\substack{p \leq P \\ p \nmid Q}} \beta_{(p,Q)}(n) \cdot \prod_{q|Q} \beta_{(q,Q)}(n) = \\
& \sum_{b \geq 0} \prod_{\substack{p \leq P \\ n^2 \equiv 4 \pmod{p^{2b}} \\ (n^2-4)2^{-2b} \text{ is} \\ \text{a discriminant}}} \frac{1}{p^b} \left(1 - \frac{1}{p} \chi_{(n^2-4)p^{-2b}}(p) \right)^{-1} \times \\
& \times \prod_{q|Q} \left\{ 1 + \left(\frac{2}{(n^2-4)q^{-2ord_q v}} \right) \begin{array}{l} , \quad q^2 \mid (n^2-4)q^{-2ord_q v} \\ , \quad q^2 \nmid (n^2-4)q^{-2ord_q v} \end{array} \right\} = \\
& \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2-4 \\ p|v \implies p \leq P}} \left[\frac{1}{v} \prod_{p \leq P} \left(1 - \frac{1}{p} \chi_D(p) \right)^{-1} \right] \times \\
& \times \prod_{q|Q} \left\{ 1 + \left(\frac{2}{(n^2-4)q^{-2ord_q v}} \right) \begin{array}{l} , \quad q^2 \mid (n^2-4)q^{-2ord_q v} \\ , \quad q^2 \nmid (n^2-4)q^{-2ord_q v} \end{array} \right\} = \beta_{P,Q}(n). \quad \blacksquare
\end{aligned}$$

5 Limit periodic functions and Fourier analysis

Let $s \geq 1$. For $f : \mathbb{N} \rightarrow \mathbb{C}$, define the seminorm

$$\|f\|_s := \left(\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} |f(n)|^s \right)^{1/s} \in [0, \infty).$$

A function f is called s -limit periodic if for every $\varepsilon > 0$ there is a periodic function h with $\|f - h\|_s \leq \varepsilon$. The set $\mathcal{D}^s$ of all s -limit periodic functions becomes a Banach space with norm $\|\cdot\|_s$ if functions f_1, f_2 with $\|f_1 - f_2\|_s = 0$ are

identified. If $1 \leq s_1 \leq s_2 < \infty$, we have $\mathcal{D}^1 \supseteq \mathcal{D}^{s_1} \supseteq \mathcal{D}^{s_2}$ as sets (but they are endowed with different norms). For all $f \in \mathcal{D}^1$, the mean value

$$M(f) := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} f(n)$$

exists. The space $\mathcal{D}^2$ is a Hilbert space with inner product

$$\langle f, h \rangle := M(f \bar{h}), \quad f, h \in \mathcal{D}^2.$$

For $u \in \mathbb{R}$, define $e_u(n) := e^{2\pi i u n}$, $n \in \mathbb{N}$. In $\mathcal{D}^2$, we have canonical orthonormal base $\{e_{a/b}\}$, where $1 \leq a \leq b$ and $\gcd(a, b) = 1$.

For all $f \in \mathcal{D}^1$, the Fourier coefficients $\hat{f}(u) := M(fe_{-u})$, $u \in \mathbb{R}$, exist.

Lemma 5.1 *For $f \in \mathcal{D}^1$, $u \notin \mathbb{Q}$, we have $\hat{f}(u) = 0$.*

Proof. Let $f \in \mathcal{D}^1$. For any $\varepsilon > 0$ there is a linear combination $\sum_{1 \leq v \leq V} e_v(n)$,

such that $\left\| f - \sum_{1 \leq v \leq V} e_v \right\|_1 < \varepsilon$, where $v \in \mathbb{Q}$. So $\left| f(n) - \sum_{1 \leq v \leq V} e_v(n) \right| < \varepsilon$, for all $1 \leq n \leq N$. Therefore we have

$$\begin{aligned} \left| \hat{f}(u) - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \sum_{1 \leq v \leq V} e_v(n) e_{-u}(n) \right| &= \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \left| \sum_{1 \leq n \leq N} \left(f(n) - \sum_{1 \leq v \leq V} e_v(n) \right) e_{-u}(n) \right| \leq \\ &\leq \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \varepsilon |e_{-u}(n)| \leq \varepsilon. \end{aligned}$$

Let $u \notin \mathbb{Q}$, since $v - u \notin \mathbb{Q}$ we have

$$\begin{aligned} \left| \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \sum_{1 \leq v \leq V} e_v(n) e_{-u}(n) \right| &= \left| \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \sum_{1 \leq v \leq V} e^{2\pi i n(v-u)} \right| = \\ &= \left| \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq v \leq V} \frac{1 - e^{2\pi i N(v-u)}}{1 - e^{2\pi i(v-u)}} \right| \leq \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq v \leq V} \frac{2}{const} = 0, \end{aligned}$$

and the lemma follows. ■

6 Limit periodicity of weighted multiplicities

In this section we will prove that the weighted multiplicities function $\beta_Q(n)$ is limit periodic (prop. 6.1), and as consequence, we will obtain the formula for calculating its mean square (end of the section).

Proposition 6.1 *The functions $\beta_Q(n) \in \mathcal{D}^1$, and for $1 \leq s \leq 2$,*

$$\lim_{P \rightarrow \infty} \|\beta_Q - \beta_{P,Q}\|_s = 0.$$

Write $\beta_Q(n) - \beta_{P,Q}(n) = \Delta_P^{(1)}(n) + \Delta_P^{(2)}(n)$, where

$$\Delta_P^{(1)}(n) := \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \text{ for some } p > P}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^{2, \begin{smallmatrix} q^2 | D \\ q^2 \nmid D \end{smallmatrix}} \right\} L(1, \chi_D)$$

and

$$\begin{aligned} \Delta_P^{(2)}(n) := & \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \Rightarrow p \leq P}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^{2, \begin{smallmatrix} q^2 | D \\ q^2 \nmid D \end{smallmatrix}} \right\} \times \\ & \times \left(L(1, \chi_D) - \prod_{p \leq P} \left(1 - \frac{\chi_D(p)}{p} \right)^{-1} \right). \end{aligned}$$

Lemma 6.2 *For $P \geq Q$ we have*

$$\frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_P^{(1)}(n) \right|^2 \ll \sum_{v > P} \frac{1}{v^2}.$$

Proof. Note that

$$\Delta_P^{(1)}(n) \leq \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \text{ for some } p > P}} \frac{2^{\omega(Q)}}{v} L(1, \chi_D), \quad (6)$$

where $\omega(Q)$ is the number of prime divisors of Q . Cauchy's inequality gives

$$\left| \Delta_P^{(1)}(n) \right| \leq \left(\sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \text{ for some } p > P}} \frac{2^{2\omega(Q)}}{v^2} \right)^{1/2} \left(\sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \text{ for some } p > P}} L(1, \chi_D)^2 \right)^{1/2}.$$

For $x \geq 1$, this gives

$$\sum_{2 < n \leq x} \left| \Delta_P^{(1)}(n) \right|^2 \leq \sum_{v > P} \frac{2^{2\omega(Q)}}{v^2} \sum_{\substack{2 < n \leq x \\ D, v \geq 1; Dv^2 = n^2 - 4}} L(1, \chi_D)^2.$$

M. Peter shows [10], [11] that the last sum is

$$\sum_{\substack{2 < n \leq x \\ D, v \geq 1; Dv^2 = n^2 - 4}} L(1, \chi_D)^2 \sim \text{const} \cdot x,$$

as $x \rightarrow \infty$. Therefore we have the claim of the lemma. ■

In order to estimate $\Delta_P^{(2)}(n)$ we must compare $L(1, \chi_D)$ with a partial product of its Euler products. This is done by comparing both terms with a smoothed version of the Dirichlet series for $L(1, \chi_D)$. Let $N \geq 1$. Then

$$\Delta_P^{(2)}(n) = \Delta_{P,N}^{(2,1)}(n) + \Delta_{P,N}^{(2,2)}(n) + \Delta_{P,N}^{(2,3)}(n),$$

where

$$\Delta_{P,N}^{(2,1)}(n) := \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \Rightarrow p \leq P}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 \mid D \\ q^2 \nmid D \end{matrix} \right\} \left(L(1, \chi_D) - \sum_{l \geq 1} \frac{\chi_D(l)}{l} e^{-l/N} \right),$$

$$\Delta_{P,N}^{(2,2)}(n) := \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \Rightarrow p \leq P}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 \mid D \\ q^2 \nmid D \end{matrix} \right\} \sum_{l \geq 1: p|l \text{ for some } p > P} \frac{\chi_D(l)}{l} e^{-l/N},$$

$$\Delta_{P,N}^{(2,3)}(n) := \sum_{\substack{D, v \geq 1; Dv^2 = n^2 - 4 \\ p|v \Rightarrow p \leq P}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^2 \begin{matrix} q^2 \mid D \\ q^2 \nmid D \end{matrix} \right\} \sum_{l \geq 1: p|l \Rightarrow p \leq P} \frac{\chi_D(l)}{l} (e^{-l/N} - 1);$$

For the following approximations we will use two lemmas:

Lemma 6.3 (Sarnak)

$$\sum_{\substack{2 < n \leq x \\ d, v \geq 1 \\ dv^2 = n^2 - 4}} 1 \sim \text{const} \cdot x$$

Proof. [15, Lemma 4.2] ■

Lemma 6.4 (Peter) For $l, v \in \mathbb{N}$ and $x \geq 3$, we have

$$\sum_{\substack{2 < n \leq x \\ d \geq 1 \\ dv^2 = n^2 - 4}} \chi_d(l) \ll \frac{x}{v^{2-\varepsilon} K(l)} + v^\varepsilon l,$$

where $K(l)$ is the squarefree kernel of l and $\varepsilon > 0$ is arbitrary.

Proof. See [12], estimate (2.7). ■

Lemma 6.5 For $P \geq Q$ and $x, N \geq 1$, we have

$$\frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,3)}(n) \right|^2 \ll \left(N^{-1/2} + \sum_{l > \sqrt{N}: p|l \Rightarrow p \leq P} \frac{1}{l} \right)^2.$$

Proof. Since $|e^{-u} - 1| \ll u$ for $0 \leq u \leq 1$, we see that for $n > 2$ the inner sum in $\Delta_{P,N}^{(2,3)}(n)$ is

$$\begin{aligned} &\ll \sum_{l \geq 1: p|l \Rightarrow p \leq P} \frac{1}{l} \left| e^{-l/N} - 1 \right| \ll \sum_{l > \sqrt{N}: p|l \Rightarrow p \leq P} \frac{2}{l} + \sum_{1 < l \leq \sqrt{N}: p|l \Rightarrow p \leq P} \frac{1}{l} \frac{l}{N} \\ &\ll \sum_{l > \sqrt{N}: p|l \Rightarrow p \leq P} \frac{1}{l} + N^{-1/2} =: c_1(P, N). \end{aligned}$$

Cauchy's inequality and (6) give

$$\begin{aligned} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,3)}(n) \right|^2 &\ll \sum_{2 < n \leq x} \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} \frac{2^{\omega(Q)}}{v} \right)^2 c_1(P, N)^2 \ll \\ &\ll c_1(P, N)^2 \sum_{2 < n \leq x} \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} \frac{2^{2\omega(Q)}}{v^2} \right) \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} 1 \right) \ll \\ &\ll c_1(P, N)^2 \sum_{\substack{2 < n \leq x \\ D, v \geq 1; Dv^2 = n^2 - 4}} 1. \end{aligned}$$

By using 6.3 the result follows. ■

Lemma 6.6 For $P \geq Q$ and $x, N \geq 1$, we have

$$\frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,2)}(n) \right|^2 \ll \sum_{l > P^2} \frac{\tau(l)}{lK(l)} + \frac{1}{x^{1/3-\varepsilon}} N^2$$

where $\tau(l)$ is the number of positive divisors of l .

Proof. Let $\alpha > \frac{1}{2}$. We write

$$\begin{aligned} \Delta_{P,N}^{(2,2)}(n) &= \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v \leq n^\alpha}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^{2, \begin{smallmatrix} q^2 | D \\ q^2 \nmid D \end{smallmatrix}} \right\} \sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} + \\ &+ \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v > n^\alpha}} \frac{1}{v} \prod_{q|Q} \left\{ 1 + \left(\frac{D}{q} \right)^{2, \begin{smallmatrix} q^2 | D \\ q^2 \nmid D \end{smallmatrix}} \right\} \sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} = \\ &= \Delta_{P,N}^{(2,2,1)}(n) + \Delta_{P,N}^{(2,2,2)}(n). \end{aligned}$$

A trivial estimate gives

$$\begin{aligned} \Delta_{P,N}^{(2,2,2)}(n) &\leq \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v > n^\alpha}} \frac{2^{\omega(Q)}}{v} \sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} \ll \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v > n^\alpha}} \frac{1}{v} \sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{1}{l} e^{-l/N} \ll \\ &\log N \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v > n^\alpha}} \frac{1}{v} \ll \log N \cdot \frac{1}{n^\alpha} \tau(n^2 - 4) \ll \log N \cdot \frac{1}{n^{\alpha-\varepsilon}}. \end{aligned}$$

Thus, since $\alpha > \frac{1}{2}$, we have

$$\sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,2,2)}(n) \right|^2 \ll (\log N)^2 \cdot \sum_{2 < n \leq x} \frac{1}{n^{2(\alpha-\varepsilon)}} \ll (\log N)^2. \quad (7)$$

By Cauchy's inequality

$$\begin{aligned} \left| \Delta_{P,N}^{(2,2,1)}(n) \right| &\leq \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ v \leq n^\alpha}} \frac{2^{\omega(Q)}}{v} \left| \sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} \right| \leq \\ &\left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} \frac{2^{2\omega(Q)}}{v^2} \right)^{\frac{1}{2}} \cdot \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4; v \leq n^\alpha} \left(\sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} \right)^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Thus for $x \geq 1$,

$$\sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,2,1)}(n) \right|^2 \ll \sum_{2 < n \leq x} \sum_{D, v \geq 1; Dv^2 = n^2 - 4; v \leq x^\alpha} \left(\sum_{\substack{l \geq 1: p|l \text{ for} \\ \text{some } p > P}} \frac{\chi_D(l)}{l} e^{-l/N} \right)^2 =$$

$$= \sum_{\substack{l_1, l_2: p_i | l_i \\ \text{for some } p_i > P}} \frac{1}{l_1 l_2} e^{-(l_1 + l_2)/N} \sum_{1 \leq v \leq x^\alpha} \sum_{\substack{2 < n \leq x \\ D \geq 1 \\ Dv^2 = n^2 - 4}} \chi_D(l_1 l_2).$$

Applying Peter's lemma 6.4 to the innermost sum gives the estimate

$$\begin{aligned} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,2,1)}(n) \right|^2 &\ll \sum_{l > P^2} \frac{1}{l} \tau(l) \sum_{1 \leq v \leq x^\alpha} \frac{x}{v^{2-\varepsilon} K(l)} + \sum_{l_1, l_2 \geq 1} \frac{l_1 l_2}{l_1 l_2} e^{-(l_1 + l_2)/N} \sum_{1 \leq v \leq x^\alpha} v^\varepsilon \\ &\ll x \sum_{l > P^2} \frac{\tau(l)}{l K(l)} + N^2 x^{\alpha(1+\varepsilon)}. \end{aligned}$$

Thus together with (7), for $\alpha = 2/3 > 1/2$ we have

$$\frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,2)}(n) \right|^2 \ll \sum_{l > P^2} \frac{\tau(l)}{l K(l)} + \frac{1}{x} (\log N)^2 + \frac{1}{x^{1/3-\varepsilon}} N^2.$$

■

In order to estimate $\Delta_{P,N}^{(2,1)}(n)$ we must show that the error

$$I(D, N) := L(1, \chi_D) - \sum_{l \geq 1} \frac{\chi_D(l)}{l} e^{-l/N},$$

which comes from smoothing Dirichlet series expansion of $L(1, \chi_D)$, is small for large N .

Lemma 6.7 *For $1/2 < \sigma_0 < 1$ define the rectangle*

$$R_x := \{s \in \mathbb{C} \mid \sigma_0 \leq \operatorname{Re}(s) \leq 1, |\operatorname{Im}(s)| \leq \log^2 x\}.$$

(a) *If $L(s, \chi_D)$ has no zeros in R_x and $D \leq x^2$, then*

$$\text{for } \operatorname{Re}(s) = \kappa, |\operatorname{Im}(s)| \leq (\log x)^2/2 \quad I(D, N) \ll x^\varepsilon N^{(\kappa-1)};$$

(b) *If $L(s, \chi_D)$ has zeros in R_x , then*

$$\#\{(n, v, D) \mid 2 < n \leq x, D, v \geq 1, n^2 - Dv^2 = 4, L(s, \chi_D) \text{ has zeros in } R_x\} \ll x^{\mu+\varepsilon},$$

where $\mu := 8(1 - \sigma_0)/\sigma_0 < 1$, $\sigma_0 < \kappa < 1$

Proof. See [10, Lemma 3.6]. ■

Lemma 6.8 *There are $0 < \kappa, \mu < 1$ such that for $P \geq Q, x, N \geq 1$ and $\varepsilon > 0$ we have*

$$\frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,1)}(n) \right|^2 \ll x^\varepsilon N^{2(\kappa-1)} + x^{\mu-1+\varepsilon} (\log(x^2 N))^2.$$

Proof. Note that a trivial estimation gives $I(D, N) \ll \log(DN)$. Cauchy's inequality, previous lemma and (6) give

$$\begin{aligned} \sum_{2 < n \leq x} \left| \Delta_{P,N}^{(2,1)}(n) \right|^2 &\ll \sum_{2 < n \leq x} \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} \frac{2^{2\omega(Q)}}{v^2} \right) \left(\sum_{D, v \geq 1; Dv^2 = n^2 - 4} |I(D, N)|^2 \right) \ll \\ &\ll \sum_{\substack{2 < n \leq x; D, v \geq 1; Dv^2 = n^2 - 4 \\ L(s, \chi_D) \text{ has no zeros in } R_x}} \left(x^\varepsilon N^{(\kappa-1)} \right)^2 + \sum_{\substack{2 < n \leq x; D, v \geq 1; Dv^2 = n^2 - 4 \\ L(s, \chi_D) \text{ has a zero in } R_x}} \log^2(DN) \ll \\ &\ll x \left(x^\varepsilon N^{(\kappa-1)} \right)^2 + x^{\mu+\varepsilon} (\log(x^2 N))^2, \end{aligned}$$

which proves the lemma. ■

Now the results are collected.

Lemma 6.9 *For $P \geq Q$, we have*

$$\|\beta_Q - \beta_{P,Q}\|_2 \ll \left(\sum_{v > P} \frac{1}{v^2} \right)^{1/2} + \left(\sum_{l > P^2} \frac{\tau(l)}{lK(l)} \right)^{1/2}.$$

Proof. For $x \geq 1$ choose $N := x^{1/8}$. Then previous lemmas show that

$$\begin{aligned} \frac{1}{x} \sum_{2 < n \leq x} \left| \Delta_P^{(2)}(n) \right|^2 &\ll \left(x^{-1/16} + \sum_{l > x^{1/16}; p|l \Rightarrow p \leq P} \frac{1}{l} \right)^2 + \sum_{l > P^2} \frac{\tau(l)}{lK(l)} + \\ &+ \frac{1}{x} (\log x)^2 + \frac{1}{x^{1/12-\varepsilon}} + x^{(\kappa-1)/4+\varepsilon} + x^{\mu-1+\varepsilon} (\log x)^2. \end{aligned}$$

Since the series

$$\sum_{l \geq 1; p|l \Rightarrow p \leq P} \frac{1}{l}$$

converges, we have for $P \geq Q$ fixed

$$\left\| \Delta_P^{(2)}(n) \right\|_2^2 \ll \sum_{l > P^2} \frac{\tau(l)}{lK(l)}.$$

Together with Lemma 6.2 this proves the claim. ■

Now we are able to prove the proposition 6.1

Proof of the proposition 6.1. By previous lemma we have

$$\|\beta_Q - \beta_{P,Q}\|_2 \ll \left(\sum_{v > P} \frac{1}{v^2} \right)^{1/2} + \left(\sum_{l > P^2} \frac{\tau(l)}{lK(l)} \right)^{1/2};$$

here

$$\sum_{v > P} \frac{1}{v^2} \longrightarrow 0,$$

as $P \rightarrow \infty$, since the series $\sum_{v \geq 1} \frac{1}{v^2}$ converges. Furthermore,

$$\sum_{l > P^2} \frac{\tau(l)}{lK(l)} \longrightarrow 0,$$

as $P \rightarrow \infty$, since

$$\sum_{l \geq 1} \frac{\tau(l)}{lK(l)} = \sum_{a, b \geq 1: a \text{ squarefree}} \frac{\tau(ab^2)}{ab^2 \cdot a} \ll \sum_{a \geq 1} \frac{a^\varepsilon}{a^2} \sum_{b \geq 1} \frac{b^{2\varepsilon}}{b^2} < \infty.$$

Thus $\lim_{P \rightarrow \infty} \|\beta_Q - \beta_{P,Q}\|_2 = 0$. For $f : \mathbb{N} \rightarrow \mathbb{C}$ arbitrary and $1 \leq s \leq 2$ we have $\|f\|_s \leq \|f\|_2$ by Hölder's inequality. Thus $\lim_{P \rightarrow \infty} \|\beta_Q - \beta_{P,Q}\|_s = 0$, for all $1 \leq s \leq 2$ and, in particular $\lim_{P \rightarrow \infty} \|\beta_Q - \beta_{P,Q}\|_1 = 0$.

Since the b -th summand of $\beta_{(p,Q)}$ is p^{2b+1} -periodic for $p \nmid Q$, 2^{2b+3} -periodic in case $p = 2$, and p^{2b+2} -periodic in case $p \mid Q$, and the series representing $\beta_{(p,Q)}$ is uniformly convergent, the function $\beta_{(p,Q)}$ is uniformly limit periodic, i.e. $\beta_{(p,Q)} \in \mathcal{D}^u$; here $\mathcal{D}^u$ is the set of all functions which can be approximated to an arbitrary accuracy by periodic functions with respect to the supremum norm. Since $\mathcal{D}^u$ is closed under multiplication it follows from Lemma 4.1, that $\beta_{P,Q} \in \mathcal{D}^u$ for all $P \geq Q$. This gives $\beta_Q \in \mathcal{D}^s$ for all $1 \leq s \leq 2$. ■

So we have now

$$\widehat{\beta_Q}(0) := M(\beta_Q) := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \beta_Q(n).$$

One can prove the

Lemma 6.10 For $b \in \mathbb{N}$, $a \in \mathbb{Z}$, $\gcd(a, b) = 1$, choose $a_p \in \mathbb{Z}$ for all $p \mid b$ such that $\sum_{p \mid b} a_p p^{-\text{ord}_p b} \equiv ab^{-1} \pmod{1}$. Then

$$\widehat{\beta_Q}\left(\frac{a}{b}\right) = \prod_{p \mid b} \widehat{\beta_{(p, Q)}}\left(\frac{a_p}{p^{\text{ord}_p b}}\right). \quad (8)$$

Proof. Word by word the proof of the same fact in [10, Lemma 4.3] ■

Corollary 6.11

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{1 \leq n \leq N} \beta_Q(n) = \widehat{\beta_Q}(0) = 1$$

By Parseval's equality and by previous lemma and corollary

$$\begin{aligned} M(\beta\bar{\beta}) &:= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n) = \sum_{b \geq 1} \sum_{\substack{1 \leq a \leq b \\ \gcd(a, b) = 1}} \left| \widehat{\beta_Q}\left(\frac{a}{b}\right) \right|^2 = \\ &= \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} \sum_{\substack{1 \leq a \leq p^c \\ a \not\equiv 0 \pmod{p}}} \left| \widehat{\beta_{(p, Q)}}\left(\frac{a}{p^c}\right) \right|^2 \right). \end{aligned} \quad (9)$$

Here the term 1 in a brackets is a contribution of $c = 0$, that is $\left| \widehat{\beta_{(p, Q)}}(0) \right|^2$.

7 Calculating the mean square of weighted multiplicities $\beta_Q(n)$

In this section we will prove Theorem 1.2.

Define the functions

$$\beta_{(p, Q, b)}(n) := \left(1 - \frac{1}{p} \chi_{(n^2 - 4)p^{-2b}}(p) \right)^{-1} \cdot \mathbb{I}_{p^b}(n),$$

and calculate the Fourier coefficients of the $\beta_{(p, Q)}(n)$ by the Fourier coefficients of the $\beta_{(p, Q, b)}(n)$.

$$\widehat{\beta_{(p, Q)}}(r) = \sum_{b \geq 0} \frac{1}{p^b} \widehat{\beta_{(p, Q, b)}}(r) \quad (10)$$

In [10] was proved that for all $p \nmid Q$,

$$\widehat{\beta_{(p, Q)}}(0) = 1 \quad (11)$$

We will prove that $\widehat{\beta_{(q, Q)}}(0) = 1$ holds as well, for all $q \mid Q$.

7.1 Calculation of the period of $\beta_{(q,Q,b)}(n)$

Let us calculate now the minimal period of the function defined above:

$$\beta_{(q,Q,b)}(n) := \left(1 - \frac{1}{q} \chi_{(n^2-4)q^{-2b}}(q)\right)^{-1} \cdot \mathbb{I}_{q^b}(n)$$

Lemma 7.1 *For $q \mid Q$ the minimal period of the $\beta_{(q,Q,b)}(n)$ is q^{2b+2} .*

Proof. a) we will find a period of a χ . We will look for a minimal k , such that $\chi(n+k) = \chi(n)$ for all n . That is

$$\left(\frac{(n^2-4)q^{-2b}}{q}\right) = \left(\frac{((n+k)^2-4)q^{-2b}}{q}\right) = \left(\frac{(n^2-4)q^{-2b} + (2nk+k^2)q^{-2b}}{q}\right)$$

and it's true for $k = q^{2b+1}$.

b) we will find a period of the $\mathbb{I}_{q^b}(n)$. We will look for a minimal k , such that $\mathbb{I}_{q^b}(n+k) = \mathbb{I}_{q^b}(n)$ for all n .

$$n^2 - 4 \equiv 0 \pmod{q^{2b}} \iff (n+k)^2 - 4 \equiv 0 \pmod{q^{2b}} \iff$$

$$n^2 - 4 + 2nk + k^2 \equiv 0 \pmod{q^{2b}} \iff 2nk + k^2 \equiv 0 \pmod{q^{2b}},$$

and it's true for $k = q^{2b}$.

$$(n^2-4)q^{-2b} \equiv 0 \pmod{q^2} \iff ((n+k)^2-4)q^{-2b} \equiv 0 \pmod{q^2} \iff$$

$$(n^2-4)q^{-2b} + (2nk+k^2)q^{-2b} \equiv 0 \pmod{q^2} \iff (2nk+k^2)q^{-2b} \equiv 0 \pmod{q^2},$$

and it's true for $k = q^{2b+2}$.

So the minimal period of the $\beta_{(q,Q,b)}(n)$ is q^{2b+2} that is $\beta_{(q,Q,b)}(n+q^{2b+2}) = \beta_{(q,Q,b)}(n)$ for all n . ■

7.2 Calculation of the Fourier coefficients $\widehat{\beta_{(q,Q,b)}}(r)$ and $\widehat{\beta_{(q,Q)}}(r)$

Theorem 7.2 *For any prime $q \mid Q$ the Fourier coefficients $\widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right)$ are:*

$$\begin{aligned}
c = 0, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}(0) &= 1 - \frac{2}{q^2(q-1)} \\
c = 0, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}(0) &= \frac{2(q^2 + q + 1)}{q^{2b+2}} \\
c = 2b + 2, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{2}{q^{2b+2}} \cos\left(\frac{4\pi a}{q^c}\right) \\
c = 2b + 1, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{1}{q^{2b+2}} \left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \times \\
&\times \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right), \\
&\text{where } \epsilon_q = \begin{cases} 1, & q \equiv 1 \pmod{4} \\ i, & q \equiv 3 \pmod{4} \end{cases} \\
c \leq 2b, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{2}{q^{2b+2}} \cos\left(\frac{4\pi a}{q^c}\right) (q^2 + q + 1) \\
c = 1, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}\left(\frac{a}{q}\right) &= -\frac{2}{q^2(q-1)} \cos\left(\frac{4\pi a}{q}\right) + \\
&\frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} \\
c = 2, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}\left(\frac{a}{q^2}\right) &= \frac{2}{q^2} \cos\left(\frac{4\pi a}{q^2}\right).
\end{aligned}$$

First we calculate $\widehat{\beta_{(q,Q,b)}}(0)$.

Remark 7.3 In all of the sums below we want to be sure that the function $\beta_{(q,Q,b)}(n)$ is defined at n , that is n is a trace of some element of $\Gamma_0(Q)$. The necessary and sufficient condition for n to be a trace of some element of $\Gamma_0(Q)$ is the condition that $\left(\frac{n^2-4}{q}\right) \neq -1$ for all $q \mid Q$. We will easily see that those n , for which $\left(\frac{n^2-4}{q}\right) \neq -1$ do not contribute to the sum. That is why we can sum over all n of the given range without any restriction.

a) $b = 0$,

$$\begin{aligned}
\widehat{\beta_{(q,Q,0)}}(0) &= \frac{1}{q^2} \sum_{n(\bmod q^2)} \left(1 - \frac{1}{q} \left(\frac{n^2-4}{q}\right)\right)^{-1} \cdot \left\{ \begin{array}{ll} 2, & q^2 \mid n^2-4 \\ \left(\frac{n^2-4}{q}\right) + 1, & q^2 \nmid n^2-4 \end{array} \right\} \\
&= \frac{1}{q^2} \sum_{n(\bmod q^2)} \left(1 - \frac{1}{q} \left(\frac{n^2-4}{q}\right)\right)^{-1} \cdot \left\{ \begin{array}{ll} 2, & n = \pm 2, \text{ or } \left(\frac{n^2-4}{q}\right) = 1 \\ 1, & n \neq \pm 2, q \mid n^2-4 \end{array} \right\} \\
&= \frac{1}{q^2} (2 \cdot 2 + 2 \cdot \#\{n(\bmod q^2) \mid \left(\frac{n^2-4}{q}\right) = 1\}) \left(1 - \frac{1}{q}\right)^{-1} + \\
&\quad \#\{n \neq \pm 2(\bmod q^2) \mid q \mid n^2-4\} = \frac{1}{q^2} (2 \cdot 2 + 2 \frac{q-3}{2} \cdot q) \left(1 - \frac{1}{q}\right)^{-1} + 2(q-1) \\
&= \frac{1}{q^2} (4 + q(q-3) \frac{q}{q-1} + 2(q-1)) = 1 - \frac{2}{q^2(q-1)}.
\end{aligned}$$

b) $b \neq 0$,

$$\begin{aligned}
\widehat{\beta_{(q,Q,b)}}(0) &= \frac{1}{q^{2b+2}} \sum_{n(\bmod q^{2b+2})} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \times \\
&\quad \times \left\{ \begin{array}{lll} 2, & n^2 = 4(\bmod q^{2b}), & q^2 \mid (n^2-4)q^{-2b} \\ 1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right), & n^2 = 4(\bmod q^{2b}), & q^2 \nmid (n^2-4)q^{-2b} \\ 0, & \text{else} \end{array} \right\} = \\
&= \frac{1}{q^{2b+2}} (2 \cdot 2 + 2 \left(1 - \frac{1}{q}\right)^{-1} \#\left\{n(\bmod q^{2b+2}) \mid \left\{ \begin{array}{l} n^2 \equiv 4(\bmod q^{2b}) \\ \left(\frac{(n^2-4)q^{-2b}}{q}\right) = 1 \end{array} \right\} \right\} + \\
&\quad \#\left\{n \neq \pm 2(\bmod q^{2b+2}) \mid \left\{ \begin{array}{l} n^2 \equiv 4(\bmod q^{2b}) \\ q^{2b+1} \mid n^2-4 \end{array} \right\} \right\}).
\end{aligned}$$

Lemma 7.4 *The cardinality of the set*

$$\left\{ n(\bmod q^{2b+2}) \mid \left\{ \begin{array}{l} n^2 \equiv 4(\bmod q^{2b}) \\ \left(\frac{(n^2-4)q^{-2b}}{q}\right) = 1 \end{array} \right\} \right\} \text{ is } q(q-1),$$

and the cardinality of the set

$$\left\{ n \neq \pm 2(\bmod q^{2b+2}) \mid \left\{ \begin{array}{l} n^2 \equiv 4(\bmod q^{2b}) \\ q^{2b+1} \mid n^2-4 \end{array} \right\} \right\} \text{ is } 2q-2.$$

Proof. a) There are $2q^2$ numbers n modulo q^{2b+2} such that $n^2 - 4 = 0 \pmod{q^{2b}}$. They are of the form kq^{2b} , where $k = \pm 1, \pm 2, \dots, \pm q^2$. So it's need to check how many of the k 's are squares modulo q . There are $(q-1)/2$ squares modulo q , hence there are $2q(q-1)/2 = q(q-1)$ numbers in the first set.

b) The number of $n \neq \pm 2$ modulo q^{2b+2} such that $n^2 = 4$ modulo q^{2b+1} is $2q - 2$. ■

So

$$\widehat{\beta_{(q,Q,b)}}(0) = \frac{1}{q^{2b+2}}(4 + 2\frac{q}{q-1}q(q-1) + 2q - 2) = \frac{2(q^2 + q + 1)}{q^{2b+2}}.$$

From the relation (10) it follows that

$$\begin{aligned} \widehat{\beta_{(q,Q)}}(0) &= \sum_{b \geq 0} \frac{1}{q^b} \widehat{\beta_{(q,Q,b)}}(0) = 1 - \frac{2}{q^2(q-1)} + \sum_{b \geq 1} \frac{1}{q^b} \frac{2(q^2 + q + 1)}{q^{2b+2}} = \\ 1 - \frac{2}{q^2(q-1)} + \frac{2(q^2 + q + 1)}{q^2} \sum_{b \geq 1} \frac{1}{q^{3b}} &= 1 - \frac{2}{q^2(q-1)} + \frac{2(q^2 + q + 1)}{q^2} \frac{1}{q^3 - 1} = 1. \end{aligned}$$

Now we will compute the Fourier coefficients $\widehat{\beta_{(q,Q,b)}}(\frac{a}{q^c})$.

a) $b \neq 0$, $c = 2b + 2$.

We will need a lemmas before we will start .

Lemma 7.5 *If $q \nmid a$, then*

$$\sum_{k=1}^{q^2} \left(\frac{k}{q}\right) e^{-2\pi i k \frac{a}{q^2}} = 0,$$

for q prime.

Proof. Note that

$$\left(\frac{m}{q}\right) = \left(\frac{m + lq}{q}\right),$$

for $l \in \mathbb{Z}$. Now we can write

$$\sum_{k=1}^{q^2} \left(\frac{k}{q}\right) e^{-2\pi i k \frac{a}{q^2}} = \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m + lq}{q}\right) e^{-2\pi i (m+lq) \frac{a}{q^2}} =$$

$$\sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m}{q}\right) e^{-2\pi i m \frac{a}{q^2}} \cdot e^{-2\pi i l \frac{a}{q}} = \sum_{m=1}^q \left(\frac{m}{q}\right) e^{-2\pi i m \frac{a}{q^2}} \sum_{l=0}^{q-1} e^{-2\pi i l \frac{a}{q}}.$$

But

$$\sum_{l=0}^{q-1} e^{-2\pi i l \frac{a}{q}} = 0$$

and hence

$$\sum_{k=1}^{q^2} \left(\frac{k}{q}\right) e^{-2\pi i k \frac{a}{q^2}} = 0,$$

as desired. ■

Lemma 7.6 For $c = 2b + 2$

$$\sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2 - 4 \\ q^{2b} \mid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{(n^2 - 4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2 - 4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} = 0.$$

Proof.

$$\sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2 - 4 \\ q^{2b} \mid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{(n^2 - 4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2 - 4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} =$$

$$e^{-4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{k}{q}\right)\right) \left(1 - \frac{1}{q} \left(\frac{k}{q}\right)\right)^{-1} e^{-2\pi i k a q^{2b-c}} +$$

$$e^{4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{-k}{q}\right)\right) \left(1 - \frac{1}{q} \left(\frac{-k}{q}\right)\right)^{-1} e^{-2\pi i k a q^{2b-c}} = \{c = 2b + 2\} =$$

$$= e^{-4\pi i \frac{a}{q^c}} \sum_{k=1, \left(\frac{k}{q}\right)=1}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i k \frac{a}{q^2}} +$$

$$e^{4\pi i \frac{a}{q^c}} \sum_{k=1, \left(\frac{-k}{q}\right)=1}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i k \frac{a}{q^2}}$$

$$\begin{aligned}
&= e^{-4\pi i \frac{a}{q^c}} \cdot 2 \left(1 - \frac{1}{q}\right)^{-1} \cdot \frac{1}{2} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q^2}} \left(1 + \left(\frac{k}{q}\right)\right) + \\
&\quad e^{4\pi i \frac{a}{q^c}} \cdot 2 \left(1 - \frac{1}{q}\right)^{-1} \cdot \frac{1}{2} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q^2}} \left(1 + \left(\frac{-k}{q}\right)\right) \\
&= e^{-4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(\frac{k}{q}\right) e^{-2\pi i k \frac{a}{q^2}} + e^{-4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q^2}} + \\
&\quad e^{4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(\frac{-k}{q}\right) e^{-2\pi i k \frac{a}{q^2}} + e^{4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q^2}}.
\end{aligned}$$

By the previous lemma and the fact that

$$\sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q^2}} = 0$$

the expression we want to compute is equal to 0. ■

Now it will be much easier to compute what we want to compute:

$$\begin{aligned}
&\widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) = \frac{1}{q^{2b+2}} \sum_{n \pmod{q^{2b+2}}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \times \\
&\times \left\{ \begin{array}{ll} \frac{2}{q}, & n^2 = 4 \pmod{q^{2b}}, \quad q^2 \mid (n^2-4)q^{-2b} \\ 1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right), & n^2 = 4 \pmod{q^{2b}}, \quad q^2 \nmid (n^2-4)q^{-2b} \\ 0, & \text{else} \end{array} \right\} \cdot e^{-2\pi i n \frac{a}{q^c}} \\
&= \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + \right. \\
&\quad \left. \sum_{\substack{n \pmod{q^{2b+2}} \\ n^2=4 \pmod{q^{2b}} \\ q^2 \nmid (n^2-4)q^{-2b}}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + \sum_{\substack{n \not\equiv \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} | n^2 - 4}} e^{-2\pi i n \frac{a}{q^c}} + \right. \\
&\quad \left. \sum_{\substack{n \not\equiv \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} | n^2 - 4 \\ q^{2b} | n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{(n^2 - 4)q^{-2b}}{q} \right) \right)^{-1} \left(1 + \left(\frac{(n^2 - 4)q^{-2b}}{q} \right) \right) e^{-2\pi i n \frac{a}{q^c}} \right) = \\
&\quad \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + 2 \cos \left(\frac{4\pi a}{q^c} \right) \sum_{k=1}^{q-1} e^{-2\pi i k a q^{2b-c+1}} + 0 \right).
\end{aligned}$$

by the previous lemma. Note that

$$\sum_{k=1}^{q-1} e^{-2\pi i k a q^{2b-c+1}} = -1$$

when $c = 2b + 2$. Therefore

$$\begin{aligned}
\widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{1}{q^{2b+2}} \left(2e^{-4\pi i \frac{a}{q^c}} + 2e^{4\pi i \frac{a}{q^c}} + 2 \cos \left(\frac{4\pi a}{q^c} \right) \cdot (-1) \right) = \\
&= \frac{1}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right) (4 - 2) = \frac{2}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right).
\end{aligned}$$

b) $b \neq 0$, $c = 2b + 1$.

The same way as before we will need two lemmas:

Lemma 7.7

$$\sum_{k=1}^{q^2} \left(\frac{k}{q} \right) e^{-2\pi i k \frac{a}{q}} = q^{\frac{3}{2}} \left(\frac{-a}{q} \right) \epsilon_q, \text{ where } \epsilon_q = \begin{cases} 1, & q \equiv 1 \pmod{4} \\ i, & q \equiv 3 \pmod{4} \end{cases}.$$

Proof.

$$\begin{aligned}
\sum_{k=1}^{q^2} \left(\frac{k}{q} \right) e^{-2\pi i k \frac{a}{q}} &= \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m+lq}{q} \right) e^{-2\pi i (m+lq) \frac{a}{q}} = \\
&= \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m}{q} \right) e^{-2\pi i m \frac{a}{q}} \cdot e^{-2\pi i l a} = q \sum_{m=1}^q \left(\frac{m}{q} \right) e^{-2\pi i m \frac{a}{q}} = \\
&= q \left(\frac{-1}{q} \right) \sum_{m=1}^q \left(\frac{-m}{q} \right) e^{2\pi i (-m) \frac{a}{q}} = q \left(\frac{-1}{q} \right) \sum_{m=1}^q \left(\frac{m}{q} \right) e^{2\pi i m \frac{a}{q}} = \\
&= q \left(\frac{-1}{q} \right) \left(\frac{a}{q} \right) \sum_{m=1}^q \left(\frac{m}{q} \right) e^{2\pi i \frac{m}{q}}.
\end{aligned}$$

The equality holds by using of property of the Gaussian sum. The value of the last sum is $\epsilon_q \sqrt{q}$, where ϵ_q defined as above. For this see, for example, [7, chapter 6]. Hence

$$\sum_{k=1}^{q^2} \left(\frac{k}{q} \right) e^{-2\pi i k \frac{a}{q}} = q \left(\frac{-1}{q} \right) \left(\frac{a}{q} \right) \epsilon_q \sqrt{q} = q^{\frac{3}{2}} \left(\frac{-a}{q} \right) \epsilon_q,$$

and we have the claim of the lemma. ■

Lemma 7.8 For $c = 2b + 1$, and ϵ_q defined as before

$$\begin{aligned}
&\sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right) e^{-2\pi i n \frac{a}{q^c}} = \\
&= \left(1 - \frac{1}{q} \right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q} \right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q} \right) \right] - 2q \left(1 - \frac{1}{q} \right)^{-1} \cos \left(\frac{4\pi a}{q^c} \right).
\end{aligned}$$

Proof.

$$\begin{aligned}
&\sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right) e^{-2\pi i n \frac{a}{q^c}} = \\
&= e^{-4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{k}{q} \right) \right) \left(1 - \frac{1}{q} \left(\frac{k}{q} \right) \right)^{-1} e^{-2\pi i k a q^{2b-c}} +
\end{aligned}$$

$$\begin{aligned}
& e^{4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{-k}{q}\right)\right) \left(1 - \frac{1}{q} \left(\frac{-k}{q}\right)\right)^{-1} e^{-2\pi i k a q^{2b-c}} = \{c = 2b + 1\} = \\
& = e^{-4\pi i \frac{a}{q^c}} \sum_{\substack{k=1, (\frac{k}{q})=1}}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i k \frac{a}{q}} + \\
& e^{4\pi i \frac{a}{q^c}} \sum_{\substack{k=1, (\frac{-k}{q})=1}}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i k \frac{a}{q}} = \\
& = e^{-4\pi i \frac{a}{q^c}} \cdot 2 \left(1 - \frac{1}{q}\right)^{-1} \cdot \frac{1}{2} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q}} \left(1 + \left(\frac{k}{q}\right)\right) + \\
& e^{4\pi i \frac{a}{q^c}} \cdot 2 \left(1 - \frac{1}{q}\right)^{-1} \cdot \frac{1}{2} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q}} \left(1 + \left(\frac{-k}{q}\right)\right) = \\
& = e^{-4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(\frac{k}{q}\right) e^{-2\pi i k \frac{a}{q}} + e^{-4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q}} + \\
& e^{4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(\frac{-k}{q}\right) e^{-2\pi i k \frac{a}{q}} + e^{4\pi i \frac{a}{q^c}} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q}}.
\end{aligned}$$

By the previous lemma and the fact that

$$\sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} e^{-2\pi i k \frac{a}{q}} = -q$$

we can write that our expression is equal to

$$\left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - 2q \left(1 - \frac{1}{q}\right)^{-1} \cos\left(\frac{4\pi a}{q^c}\right),$$

proving the claim. ■

By using the same arguments as in the case a) we will get

$$\begin{aligned}
\widehat{\beta_{(q,Q,b)}\left(\frac{a}{q^c}\right)} &= \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + \sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4}} e^{-2\pi i n \frac{a}{q^c}} + \right. \\
&\quad \left. \sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} \right) = \\
&= \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + 2 \cos\left(\frac{4\pi a}{q^c}\right) \sum_{k=1}^{q-1} e^{-2\pi i k a} + \right. \\
&\quad \left. \left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - 2q \left(1 - \frac{1}{q}\right)^{-1} \cos\left(\frac{4\pi a}{q^c}\right) \right).
\end{aligned}$$

by using previous lemma. Note that

$$\sum_{k=1}^{q-1} e^{-2\pi i k a} = q - 1,$$

so

$$\begin{aligned}
\widehat{\beta_{(q,Q,b)}\left(\frac{a}{q^c}\right)} &= \frac{1}{q^{2b+2}} \left(4 \cos\left(\frac{4\pi a}{q^c}\right) + 2(q-1) \cos\left(\frac{4\pi a}{q^c}\right) + \right. \\
&\quad \left. \left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - 2q \left(1 - \frac{1}{q}\right)^{-1} \cos\left(\frac{4\pi a}{q^c}\right) \right) \\
&= \frac{1}{q^{2b+2}} \left(\left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) \right).
\end{aligned}$$

c) $b \neq 0, 2b \geq c$.

By doing the same steps as before we can write

Lemma 7.9 For $2b \geq c$,

$$\begin{aligned} \sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} = \\ = 2q^2 \cos\left(\frac{4\pi a}{q^c}\right). \end{aligned}$$

Proof.

$$\begin{aligned} \sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q}\right)\right) e^{-2\pi i n \frac{a}{q^c}} = \\ = e^{-4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{k}{q}\right)\right) \left(1 - \frac{1}{q} \left(\frac{k}{q}\right)\right)^{-1} e^{-2\pi i k a q^{2b-c}} + \\ e^{4\pi i \frac{a}{q^c}} \sum_{\substack{k=1 \\ q \nmid k}}^{q^2-1} \left(1 + \left(\frac{-k}{q}\right)\right) \left(1 - \frac{1}{q} \left(\frac{-k}{q}\right)\right)^{-1} e^{-2\pi i k a q^{2b-c}} = \{2b \geq c\} = \\ = e^{-4\pi i \frac{a}{q^c}} \sum_{k=1, \left(\frac{k}{q}\right)=1}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} + e^{4\pi i \frac{a}{q^c}} \sum_{k=1, \left(\frac{-k}{q}\right)=1}^{q^2-1} 2 \left(1 - \frac{1}{q}\right)^{-1} = \\ = 2 \frac{q}{q-1} \#\{k \pmod{q^2} \mid \left(\frac{k}{q}\right) = 1\} 2 \cos\left(\frac{4\pi a}{q^c}\right) = 4 \frac{q}{q-1} \frac{q(q-1)}{2} \cos\left(\frac{4\pi a}{q^c}\right) = \\ = 2q^2 \cos\left(\frac{4\pi a}{q^c}\right). \end{aligned}$$

■

And gathering all together we have

$$\widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) = \frac{1}{q^{2b+2}} \left(\sum_{\substack{n=2 \\ n=-2+q^{2b+2}}} 2e^{-2\pi i n \frac{a}{q^c}} + 2 \cos\left(\frac{4\pi a}{q^c}\right) \sum_{k=1}^{q-1} e^{-2\pi i k a q^{2b-c+1}} + \right.$$

$$\begin{aligned}
& \sum_{\substack{n \neq \pm 2 \pmod{q^{2b+2}} \\ q^{2b+1} \nmid n^2-4 \\ q^{2b} \mid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right)^{-1} \left(1 + \left(\frac{(n^2-4)q^{-2b}}{q} \right) \right) e^{-2\pi i n \frac{a}{q^c}} \Bigg) \\
&= \frac{1}{q^{2b+2}} \left(4 \cos \left(\frac{4\pi a}{q^c} \right) + 2(q-1) \cos \left(\frac{4\pi a}{q^c} \right) + 2q^2 \cos \left(\frac{4\pi a}{q^c} \right) \right) \\
&= \frac{1}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right) [4 + 2(q-1) + 2q^2] = \frac{2}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right) (q^2 + q + 1).
\end{aligned}$$

d) $b = 0, c = 1$.

As always we need two lemmas:

Lemma 7.10

$$\sum_{n \pmod{q^2}} \left(\frac{n^2-4}{q} \right) e^{-2\pi i n \frac{a}{q}} = q \sum_{n \pmod{q}} \left(\frac{n^2-4}{q} \right) e^{-2\pi i n \frac{a}{q}}.$$

Proof.

$$\begin{aligned}
\sum_{n \pmod{q^2}} \left(\frac{n^2-4}{q} \right) e^{-2\pi i n \frac{a}{q}} &= \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{(m+lq)^2-4}{q} \right) e^{-2\pi i (m+lq) \frac{a}{q}} = \\
&= \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m^2-4}{q} \right) e^{-2\pi i m \frac{a}{q}} = q \sum_{m=1}^q \left(\frac{m^2-4}{q} \right) e^{-2\pi i m \frac{a}{q}} = \\
&= q \sum_{n \pmod{q}} \left(\frac{n^2-4}{q} \right) e^{-2\pi i n \frac{a}{q}}.
\end{aligned}$$

■

Lemma 7.11

$$\begin{aligned}
& \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q \nmid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{n^2-4}{q} \right) \right)^{-1} \left(\left(\frac{n^2-4}{q} \right) + 1 \right) e^{-2\pi i n \frac{a}{q}} = \\
&= q \left(1 - \frac{1}{q} \right)^{-1} \left[-2 \cos \left(\frac{4\pi a}{q} \right) + \sum_{n \pmod{q}} \left(\frac{n^2-4}{q} \right) e^{-2\pi i n \frac{a}{q}} \right].
\end{aligned}$$

Proof.

$$\begin{aligned}
& \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q \nmid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q}} = \\
&= \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q}} = \\
&= \sum_{\substack{n \pmod{q^2} \\ \left(\frac{n^2 - 4}{q}\right) = 1}} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i n \frac{a}{q}} = \\
&= 2 \cdot \frac{1}{2} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q}} = \\
&= \left(1 - \frac{1}{q}\right)^{-1} \left[\sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} e^{-2\pi i n \frac{a}{q}} + \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q}} \right] = \\
&= \left(1 - \frac{1}{q}\right)^{-1} \left(-2q \cos\left(\frac{4\pi a}{q}\right) + \sum_{n \pmod{q^2}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q}} - \right. \\
&\quad \left. \sum_{\substack{n \pmod{q^2} \\ n = \pm 2 \pmod{q}}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q}} \right) \\
&= \left(1 - \frac{1}{q}\right)^{-1} \left(-2q \cos\left(\frac{4\pi a}{q}\right) + \sum_{n \pmod{q^2}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q}} - 0 \right) = \\
&= \left(1 - \frac{1}{q}\right)^{-1} \left[-2q \cos\left(\frac{4\pi a}{q}\right) + q \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q}} \right].
\end{aligned}$$

by previous lemma, and that's it. ■

Now we can gather the results

$$\widehat{\beta_{(q, Q, 0)}}\left(\frac{a}{q}\right) = \frac{1}{q^2} \sum_{n \pmod{q^2}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \cdot \left\{ \begin{array}{ll} 2, & q^2 \mid n^2 - 4 \\ \left(\frac{n^2 - 4}{q}\right) + 1, & q^2 \nmid n^2 - 4 \end{array} \right\} \cdot e^{-2\pi i n \frac{a}{q}} =$$

$$\frac{1}{q^2} \left(\sum_{\substack{n=2 \\ n=-2+q^2}} 2e^{-2\pi i n \frac{a}{q}} + \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q | n^2 - 4}} e^{-2\pi i n \frac{a}{q}} + \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q \nmid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q} \right) \right)^{-1} \left(\left(\frac{n^2 - 4}{q} \right) + 1 \right) e^{-2\pi i n \frac{a}{q}} \right) =$$

$$\begin{aligned} & \frac{1}{q^2} \left(4 \cos \left(\frac{4\pi a}{q} \right) + 2(q-1) \cos \left(\frac{4\pi a}{q} \right) + \right. \\ & \quad \left. q \left(1 - \frac{1}{q} \right)^{-1} \left[-2 \cos \left(\frac{4\pi a}{q} \right) + \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right) e^{-2\pi i n \frac{a}{q}} \right] \right) = \\ & -\frac{2}{q^2(q-1)} \cos \left(\frac{4\pi a}{q} \right) + \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right) e^{-2\pi i n \frac{a}{q}}. \end{aligned}$$

e) $b = 0, c = 2$.

Lemma 7.12

$$\sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q | n^2 - 4}} e^{-2\pi i n \frac{a}{q^2}} = -2 \cos \left(\frac{4\pi a}{q^2} \right).$$

Proof.

The condition $\{ n \neq \pm 2 \pmod{q^2}, q | n^2 - 4 \}$ we can rewrite in the form $\{ n = \pm 2 + kq, k = 1, \dots, q-1 \}$. Thus

$$\begin{aligned} & \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q | n^2 - 4}} e^{-2\pi i n \frac{a}{q^2}} = \sum_{k=1}^{q-1} e^{-2\pi i (\pm 2 + kq) \frac{a}{q^2}} = \\ & = 2 \cos \left(\frac{4\pi a}{q^2} \right) \sum_{k=1}^{q-1} e^{-2\pi i k \frac{a}{q}} = -2 \cos \left(\frac{4\pi a}{q^2} \right), \end{aligned}$$

and we are done. ■

Lemma 7.13

$$\sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q \nmid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q^2}} = 0.$$

Proof.

$$\begin{aligned} & \sum_{\substack{n \neq \pm 2 \pmod{q^2} \\ q \nmid n^2 - 4}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q^2}} = \\ &= \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(1 - \frac{1}{q} \left(\frac{n^2 - 4}{q}\right)\right)^{-1} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q^2}} = \\ &= \sum_{\substack{n \pmod{q^2} \\ \left(\frac{n^2 - 4}{q}\right) = 1}} 2 \left(1 - \frac{1}{q}\right)^{-1} e^{-2\pi i n \frac{a}{q^2}} = \\ &= 2 \cdot \frac{1}{2} \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(\left(\frac{n^2 - 4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q^2}} = \\ &= \left(1 - \frac{1}{q}\right)^{-1} \left[\sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} e^{-2\pi i n \frac{a}{q^2}} + \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q^2}} \right]. \end{aligned}$$

The first sum in the brackets is 0, and so we need to compute an expression

$$\begin{aligned} & \left(1 - \frac{1}{q}\right)^{-1} \sum_{\substack{n \pmod{q^2} \\ n \neq \pm 2 \pmod{q}}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q^2}} = \\ &= \left(1 - \frac{1}{q}\right)^{-1} \sum_{n \pmod{q^2}} \left(\frac{n^2 - 4}{q}\right) e^{-2\pi i n \frac{a}{q^2}} = \\ &= \left(1 - \frac{1}{q}\right)^{-1} \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{(m + lq)^2 - 4}{q}\right) e^{-2\pi i (m + lq) \frac{a}{q^2}} = \\ &= \left(1 - \frac{1}{q}\right)^{-1} \sum_{m=1}^q \sum_{l=0}^{q-1} \left(\frac{m^2 - 4}{q}\right) e^{-2\pi i m \frac{a}{q^2}} \cdot e^{-2\pi i l \frac{a}{q}} = \end{aligned}$$

$$= \left(1 - \frac{1}{q}\right)^{-1} \sum_{m=1}^q \left(\frac{m^2-4}{q}\right) e^{-2\pi i m \frac{a}{q^2}} \sum_{l=0}^{q-1} e^{-2\pi i l \frac{a}{q}} = 0,$$

since the last sum is 0. ■

$$\begin{aligned} \widehat{\beta_{(q,Q,0)}}\left(\frac{a}{q^2}\right) &= \frac{1}{q^2} \sum_{n(\bmod q^2)} \left(1 - \frac{1}{q} \left(\frac{n^2-4}{q}\right)\right)^{-1} \cdot \left\{ \begin{matrix} 2, & q^2 \mid n^2-4 \\ \left(\frac{n^2-4}{q}\right) + 1, & q^2 \nmid n^2-4 \end{matrix} \right\} e^{-2\pi i n \frac{a}{q^2}} = \\ &= \frac{1}{q^2} \left(\sum_{\substack{n=2 \\ n=-2+q^2}} 2e^{-2\pi i n \frac{a}{q^2}} + \sum_{\substack{n \neq \pm 2(\bmod q^2) \\ q \nmid n^2-4}} e^{-2\pi i n \frac{a}{q^2}} + \right. \\ &\quad \left. \sum_{\substack{n \neq \pm 2(\bmod q^2) \\ q \nmid n^2-4}} \left(1 - \frac{1}{q} \left(\frac{n^2-4}{q}\right)\right)^{-1} \left(\left(\frac{n^2-4}{q}\right) + 1\right) e^{-2\pi i n \frac{a}{q^2}} \right) = \\ &= \frac{1}{q^2} \left(4 \cos\left(\frac{4\pi a}{q^2}\right) - 2 \cos\left(\frac{4\pi a}{q^2}\right) + 0 \right) = \frac{2}{q^2} \cos\left(\frac{4\pi a}{q^2}\right) \end{aligned}$$

Summarize the above results:

$$c = 0, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}(0) = 1 - \frac{2}{q^2(q-1)}$$

$$c = 0, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}(0) = \frac{2(q^2 + q + 1)}{q^{2b+2}}$$

$$c = 2b + 2, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) = \frac{2}{q^{2b+2}} \cos\left(\frac{4\pi a}{q^c}\right)$$

$$\begin{aligned} c = 2b + 1, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{1}{q^{2b+2}} \left(\left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \times \right. \\ &\quad \left. \times \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) \right) \end{aligned}$$

$$\begin{aligned}
c \leq 2b, \quad b \neq 0, \quad \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q^c}\right) &= \frac{2}{q^{2b+2}} \cos\left(\frac{4\pi a}{q^c}\right) (q^2 + q + 1) \\
c = 1, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}\left(\frac{a}{q}\right) &= -\frac{2}{q^2(q-1)} \cos\left(\frac{4\pi a}{q}\right) + \\
&\quad \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} \\
c = 2, \quad b = 0, \quad \widehat{\beta_{(q,Q,0)}}\left(\frac{a}{q^2}\right) &= \frac{2}{q^2} \cos\left(\frac{4\pi a}{q^2}\right).
\end{aligned}$$

The theorem is completely proved.

Now we can compute the $\widehat{\beta_{(q,Q)}}(r)$.

Theorem 7.14 *The Fourier coefficients $\widehat{\beta_{(q,Q)}}\left(\frac{a}{q^c}\right)$ are:*

$$\begin{aligned}
\widehat{\beta_{(q,Q)}}\left(\frac{a}{q}\right) &= \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} \\
\widehat{\beta_{(q,Q)}}\left(\frac{a}{q^2}\right) &= \frac{2}{q(q-1)} \cos\left(\frac{4\pi a}{q^2}\right) \\
\widehat{\beta_{(q,Q)}}\left(\frac{a}{q^c}\right) &= \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) \frac{1}{q^{\frac{3c-4}{2}}}, \text{ for } c > 2, c \text{ even} \\
\widehat{\beta_{(q,Q)}}\left(\frac{a}{q^c}\right) &= \frac{1}{q-1} \frac{1}{q^{\frac{3c-4}{2}}} \epsilon_q\left(\frac{a}{q}\right) \left[e^{-4\pi i \frac{a}{q^c} \left(\frac{-1}{q}\right)} + e^{4\pi i \frac{a}{q^c}} \right], \text{ for } c > 2, c \text{ odd}
\end{aligned}$$

Proof. By using (10) we get

a) for $c = 1$,

$$\begin{aligned}
\widehat{\beta_{(q,Q)}}\left(\frac{a}{q}\right) &= \sum_{b \geq 0} \frac{1}{q^b} \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q}\right) = -\frac{2}{q^2(q-1)} \cos\left(\frac{4\pi a}{q}\right) + \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} + \\
&\quad \sum_{b \geq 1} \frac{1}{q^b} \widehat{\beta_{(q,Q,b)}}\left(\frac{a}{q}\right) = -\frac{2}{q^2(q-1)} \cos\left(\frac{4\pi a}{q}\right) + \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} + \\
&\quad \sum_{b \geq 1} \frac{1}{q^b} \frac{2}{q^{2b+2}} \cos\left(\frac{4\pi a}{q}\right) (q^2 + q + 1) = -\frac{2}{q^2(q-1)} \cos\left(\frac{4\pi a}{q}\right) + \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2-4}{q}\right) e^{-2\pi i n \frac{a}{q}} +
\end{aligned}$$

$$2 \cos \left(\frac{4\pi a}{q} \right) (q^2 + q + 1) \frac{1}{q^2(q^3 - 1)} = \frac{1}{q - 1} \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right) e^{-2\pi i n \frac{a}{q}}.$$

b) for $c = 2$,

$$\begin{aligned} \widehat{\beta_{(q,Q)}} \left(\frac{a}{q^2} \right) &= \sum_{b \geq 0} \frac{1}{q^b} \widehat{\beta_{(q,Q,b)}} \left(\frac{a}{q^2} \right) = \frac{2}{q^2} \cos \left(\frac{4\pi a}{q^2} \right) + \sum_{b \geq 1} \frac{1}{q^b} \widehat{\beta_{(q,Q,b)}} \left(\frac{a}{q^2} \right) = \\ &= \frac{2}{q^2} \cos \left(\frac{4\pi a}{q^2} \right) + \sum_{b \geq 1} \frac{1}{q^b} \frac{2}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^2} \right) (q^2 + q + 1) = \\ &= \frac{2}{q^2} \cos \left(\frac{4\pi a}{q^2} \right) + 2 \cos \left(\frac{4\pi a}{q^2} \right) (q^2 + q + 1) \frac{1}{q^2(q^3 - 1)} = \frac{2}{q(q - 1)} \cos \left(\frac{4\pi a}{q^2} \right). \end{aligned}$$

c) for $c > 2$,

$$\begin{aligned} \widehat{\beta_{(q,Q)}} \left(\frac{a}{q^c} \right) &= \sum_{0 \leq b < \frac{c-2}{2}} \frac{1}{q^b} \cdot 0 + \sum_{b = \frac{c-2}{2}} \frac{1}{q^b} \frac{2}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right) + \\ &\sum_{b = \frac{c-1}{2}} \frac{1}{q^b} \left[\frac{1}{q^{2b+2}} \left(\left(1 - \frac{1}{q} \right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q} \right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q} \right) \right] - \frac{2}{q-1} \cos \left(\frac{4\pi a}{q^c} \right) \right) \right] + \\ &\sum_{b \geq \frac{c}{2}} \frac{1}{q^b} \frac{2}{q^{2b+2}} \cos \left(\frac{4\pi a}{q^c} \right) (q^2 + q + 1). \end{aligned}$$

For c even we have

$$\begin{aligned} \widehat{\beta_{(q,Q)}} \left(\frac{a}{q^c} \right) &= \frac{1}{q^{\frac{c-2}{2}}} \frac{2}{q^c} \cos \left(\frac{4\pi a}{q^c} \right) + \frac{2}{q^2} \cos \left(\frac{4\pi a}{q^c} \right) (q^2 + q + 1) \sum_{b \geq \frac{c}{2}} \frac{1}{q^{3b}} = \\ &= 2 \cos \left(\frac{4\pi a}{q^c} \right) \left[\frac{1}{q^{\frac{3c-2}{2}}} + \frac{1}{q^2} (q^2 + q + 1) \frac{1}{q^{\frac{3c-6}{2}}} \frac{1}{q^3 - 1} \right] = \frac{2}{q - 1} \cos \left(\frac{4\pi a}{q^c} \right) \frac{1}{q^{\frac{3c-4}{2}}}. \end{aligned}$$

And for c odd

$$\frac{1}{q^{\frac{c-1}{2}}} \left[\frac{1}{q^{c+1}} \left(\left(1 - \frac{1}{q} \right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q} \right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q} \right) \right] - \frac{2}{q-1} \cos \left(\frac{4\pi a}{q^c} \right) \right) \right] +$$

$$\begin{aligned}
& \frac{2}{q^2} \cos\left(\frac{4\pi a}{q^c}\right) (q^2 + q + 1) \sum_{b \geq \frac{c+1}{2}} \frac{1}{q^{3b}} = \\
& = \frac{1}{q^{\frac{3c+1}{2}}} \left(1 - \frac{1}{q}\right)^{-1} q^{\frac{3}{2}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - \frac{1}{q^{\frac{3c+1}{2}}} \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) + \\
& \frac{2}{q^2} \cos\left(\frac{4\pi a}{q^c}\right) (q^2 + q + 1) \frac{1}{q^{\frac{3c-3}{2}}} \frac{1}{q^3 - 1} = \\
& = \frac{1}{q^{\frac{3c-4}{2}}} \frac{1}{q-1} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-a}{q}\right) + e^{4\pi i \frac{a}{q^c}} \left(\frac{a}{q}\right) \right] - \frac{1}{q^{\frac{3c+1}{2}}} \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) + \frac{1}{q^{\frac{3c+1}{2}}} \frac{2}{q-1} \cos\left(\frac{4\pi a}{q^c}\right) = \\
& = \frac{1}{q-1} \frac{1}{q^{\frac{3c-4}{2}}} \epsilon_q \left(\frac{a}{q}\right) \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-1}{q}\right) + e^{4\pi i \frac{a}{q^c}} \right]. \blacksquare
\end{aligned}$$

7.3 Calculating the mean square of weighted multiplicities function

In this subsection we will calculate the mean-square of the weighted multiplicities $\beta_Q(n)$.

To calculate the limit

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n)$$

we will use (9). Let us define the function

$$A_Q(p^c) := \sum_{\substack{1 \leq a \leq p^c \\ p \nmid a}} \left| \widehat{\beta_{(p,Q)}}\left(\frac{a}{p^c}\right) \right|^2. \quad (12)$$

So we have that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n) = \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} A_Q(p^c) \right).$$

The values of the $A_Q(p^c)$ for $p \nmid Q$ were calculate by M.Peter [10]. They are:

$$p \neq 2, p \nmid Q; \quad A_Q(p) = \frac{p^2 - 2p - 1}{(p^2 - 1)^2}, \quad A_Q(p^c) = \frac{2(p - 1)}{(p^2 - 1)^2 p^{2c-3}}$$

$$p = 2; \quad A_Q(2) = \frac{1}{9}, \quad A_Q(4) = \frac{1}{18},$$

$$A_Q(8) = 0, \quad A_Q(16) = \frac{1}{9 \cdot 16},$$

$$A_Q(32) = 0, \quad A_Q(2^c) = \frac{1}{9 \cdot 2^{2c-5}}, c \geq 6.$$

We just need to complete his work by adding the case $q \mid Q$.

a) $c = 1$;

$$\begin{aligned} A_Q(q) &= \sum_{\substack{1 \leq a \leq q \\ q \nmid a}} \left| \widehat{\beta_{(q,Q)}} \left(\frac{a}{q} \right) \right|^2 = \sum_{\substack{1 \leq a \leq q \\ q \nmid a}} \left| \frac{1}{q-1} \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right) e^{-2\pi i n \frac{a}{q}} \right|^2 = \\ &= \frac{1}{(q-1)^2} \sum_{\substack{1 \leq a \leq q \\ q \nmid a}} \sum_{n_1, n_2 \pmod{q}} \left(\frac{n_1^2 - 4}{q} \right) \left(\frac{n_2^2 - 4}{q} \right) e^{2\pi i (n_1 - n_2) \frac{a}{q}} = \\ &= \frac{1}{(q-1)^2} \sum_{n_1, n_2 \pmod{q}} \left(\frac{n_1^2 - 4}{q} \right) \left(\frac{n_2^2 - 4}{q} \right) \sum_{a=1}^{q-1} e^{2\pi i (n_1 - n_2) \frac{a}{q}}. \end{aligned}$$

The sum

$$\sum_{a=1}^{q-1} e^{2\pi i (n_1 - n_2) \frac{a}{q}} = \begin{cases} q-1, & n_1 = n_2 = n \pmod{q} \\ -1, & \text{else} \end{cases}.$$

Note that

$$\sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right) = \frac{q-3}{2} - \frac{q-1}{2} = -1,$$

hence

$$\begin{aligned} A_Q(q) &= \frac{1}{(q-1)^2} \left((q-1) \sum_{n \pmod{q}} \left(\frac{n^2 - 4}{q} \right)^2 - \sum_{n_1 \pmod{q}} \sum_{n_2 \neq n_1 \pmod{q}} \left(\frac{n_1^2 - 4}{q} \right) \left(\frac{n_2^2 - 4}{q} \right) \right) = \\ &= \frac{1}{(q-1)^2} \left((q-1)(q-2) - \sum_{n_1 \pmod{q}} \left(\sum_{n_2 \pmod{q}} \left(\frac{n_1^2 - 4}{q} \right) \left(\frac{n_2^2 - 4}{q} \right) - \left(\frac{n_1^2 - 4}{q} \right)^2 \right) \right) = \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(q-1)^2} \left((q-1)(q-2) - \left(1 - \sum_{n_1 \pmod{q}} \left(\frac{n_1^2 - 4}{q} \right)^2 \right) \right) = \\
&= \frac{1}{(q-1)^2} (q^2 - 3q + 2 - (1 - (q-2))) = \frac{q^2 - 2q - 1}{(q-1)^2}.
\end{aligned}$$

b) $c = 2$;

$$\begin{aligned}
A_Q(q^2) &= \sum_{\substack{1 \leq a \leq q^2 \\ q \nmid a}} \left| \widehat{\beta_{(q,Q)}} \left(\frac{a}{q^2} \right) \right|^2 = \sum_{\substack{1 \leq a \leq q^2 \\ q \nmid a}} \left| \frac{2}{q(q-1)} \cos \left(\frac{4\pi a}{q^2} \right) \right|^2 = \\
&= \frac{4}{q^2(q-1)^2} \sum_{\substack{1 \leq a \leq q^2 \\ q \nmid a}} \left| \cos \left(\frac{4\pi a}{q^2} \right) \right|^2 = \frac{1}{q^2(q-1)^2} \sum_{\substack{1 \leq a \leq q^2 \\ q \nmid a}} \left(e^{8\pi i \frac{a}{q^2}} + e^{-8\pi i \frac{a}{q^2}} + 2 \right) = \\
&= \frac{1}{q^2(q-1)^2} \left(2(q^2 - q) + \sum_{\substack{1 \leq a \leq q^2 \\ q \nmid a}} \left(e^{8\pi i \frac{a}{q^2}} + e^{-8\pi i \frac{a}{q^2}} \right) \right) = \frac{2(q^2 - q)}{q^2(q-1)^2} = \frac{2}{q(q-1)}.
\end{aligned}$$

c1) $c > 2$, c even;

$$\begin{aligned}
A_Q(q^c) &= \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left| \widehat{\beta_{(q,Q)}} \left(\frac{a}{q^c} \right) \right|^2 = \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left| \frac{2}{q-1} \cos \left(\frac{4\pi a}{q^c} \right) \frac{1}{q^{\frac{3c-4}{2}}} \right|^2 = \\
&= \frac{1}{q^{3c-4}} \frac{4}{(q-1)^2} \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left| \cos \left(\frac{4\pi a}{q^c} \right) \right|^2 = \frac{1}{q^{3c-4}} \frac{1}{(q-1)^2} \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left(e^{8\pi i \frac{a}{q^c}} + e^{-8\pi i \frac{a}{q^c}} + 2 \right) = \\
&= \frac{1}{q^{3c-4}} \frac{1}{(q-1)^2} \left(2(q^c - q^{c-1}) + \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left(e^{8\pi i \frac{a}{q^c}} + e^{-8\pi i \frac{a}{q^c}} \right) \right) = \frac{2q^{c-1}(q-1)}{q^{3c-4}(q-1)^2} = \frac{2}{q^{2c-3}(q-1)}.
\end{aligned}$$

c2) $c > 2$, c odd;

$$\begin{aligned}
A_Q(q^c) &= \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left| \widehat{\beta_{(q,Q)}}\left(\frac{a}{q^c}\right) \right|^2 = \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left| \frac{1}{q-1} \frac{1}{q^{\frac{3c-4}{2}}} \epsilon_q \left(\frac{a}{q} \right) \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-1}{q} \right) + e^{4\pi i \frac{a}{q^c}} \right] \right|^2 = \\
&= \frac{1}{(q-1)^2} \frac{1}{q^{3c-4}} \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-1}{q} \right) + e^{4\pi i \frac{a}{q^c}} \right] \overline{\epsilon_q \left[e^{-4\pi i \frac{a}{q^c}} \left(\frac{-1}{q} \right) + e^{4\pi i \frac{a}{q^c}} \right]} = \\
&= \frac{1}{(q-1)^2} \frac{1}{q^{3c-4}} \sum_{\substack{1 \leq a \leq q^c \\ q \nmid a}} \left(2 + \left(\frac{-1}{q} \right) \left[e^{8\pi i \frac{a}{q^c}} + e^{-8\pi i \frac{a}{q^c}} \right] \right) = \frac{2q^{c-1}(q-1)}{q^{3c-4}(q-1)^2} = \frac{2}{q^{2c-3}(q-1)}.
\end{aligned}$$

And we can see that for $c > 2$, $A_Q(q^c)$ does not depend on parity of c . Now we have by [10]

$$p \neq 2, p \nmid Q; \quad A_Q(p) = \frac{p^2 - 2p - 1}{(p^2 - 1)^2}, \quad A_Q(p^c) = \frac{2(p-1)}{(p^2 - 1)^2 p^{2c-3}}$$

$$p = 2; \quad A_Q(2) = \frac{1}{9}, \quad A_Q(4) = \frac{1}{18},$$

$$A_Q(8) = 0, \quad A_Q(16) = \frac{1}{9 \cdot 16},$$

$$A_Q(32) = 0, \quad A_Q(2^c) = \frac{1}{9 \cdot 2^{2c-5}}, c \geq 6.$$

and what we have find, for $q \mid Q$,

$$A_Q(q) = \frac{q^2 - 2q - 1}{(q-1)^2}, \quad A_Q(q^2) = \frac{2}{q(q-1)}, \quad A_Q(q^c) = \frac{2}{q^{2c-3}(q-1)}.$$

Now we can calculate

$$\begin{aligned}
\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n) &= \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} A_Q(p^c) \right) = \left(1 + \frac{1}{9} + \frac{1}{18} + \frac{1}{9 \cdot 16} + \sum_{c \geq 6} \frac{1}{9 \cdot 2^{2c-5}} \right) \times \\
&\times \prod_{q \mid Q} \left(1 + \frac{q^2 - 2q - 1}{(q-1)^2} + \frac{2}{q(q-1)} + \sum_{c \geq 2} \frac{2}{q^{2c-3}(q-1)} \right) \cdot \prod_{\substack{p \neq 2 \\ p \nmid Q}} \left(1 + \frac{p^2 - 2p - 1}{(p^2 - 1)^2} + \sum_{c \geq 2} \frac{2(p-1)}{(p^2 - 1)^2 p^{2c-3}} \right) =
\end{aligned}$$

$$= \frac{1015}{864} \prod_{q|Q} \frac{2q(q^2 - q - 1)}{(q+1)(q-1)^2} \prod_{\substack{p \neq 2 \\ p \nmid Q}} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p+1)}.$$

That is

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_Q^2(n) &= \frac{1015}{864} \prod_{q|Q} \frac{2q(q^2 - q - 1)}{(q+1)(q-1)^2} \prod_{\substack{p \neq 2 \\ p \nmid Q}} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p+1)} = \\ &= \frac{1015}{864} \prod_{q|Q} \frac{2q(q^2 - q - 1)}{(q+1)(q-1)^2} \frac{(q^2 - 1)^2(q+1)}{q^2(q^3 + q^2 - q - 3)} \prod_{p \neq 2} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p+1)} = \\ &= \frac{1015}{864} \prod_{q|Q} \frac{2(q+1)^2(q^2 - q - 1)}{q(q^3 + q^2 - q - 3)} \prod_{p \neq 2} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p+1)} \\ &= \left(\prod_{q|Q} \frac{2(q^2 - q - 1)(q+1)^2}{q(q^3 + q^2 - q - 3)} \right) \cdot 1.328... = C_1 \prod_{q|Q} \frac{2(q^2 - q - 1)(q+1)^2}{q(q^3 + q^2 - q - 3)}, \end{aligned}$$

proving the result pointed out in introduction.

8 Proof of the Theorem 1.3. Main points

Let B be an indefinite division quaternion algebra over $\mathbb{Q}$ with discriminant d_B , and R be the *maximal* order in B . Let

$$\Gamma_R = \{\alpha \in R \mid N_B(\alpha) = 1\}$$

be the unit group of R . Define the weighted multiplicities function in the similar way as for $\Gamma_0(Q)$:

$$\beta_R(n) = \frac{1}{4} \sum_{\substack{\{T\} \subset \Gamma_R \\ T=T_0^k \text{ is hyperbolic} \\ |tr T|=n}} \frac{\ln \mathcal{N}(T_0)}{\mathcal{N}(T)^{1/2} - \mathcal{N}(T)^{-1/2}}.$$

The first step is to express weighted multiplicities in the terms of Dirichlet's L -functions. The analog of the theorem 3.1 is

Theorem 8.1 *Let B be indefinite division quaternion algebra over $\mathbb{Q}$, and d_B be its reduced discriminant. Then*

$$\beta_R(n) = \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4}} \frac{1}{v} L(1, \chi_D) \prod_{p|d_B} \left\{ \begin{array}{ll} 1 - \left(\frac{D}{p}\right), & p^2 \nmid D \\ 0, & p^2 \mid D \end{array} \right\},$$

where D is a discriminant, i.e. $D \equiv 0, 1 \pmod{4}$.

Proof. One uses exactly the same techniques as for the $\Gamma_0(Q)$ case. ■

In the second step we change L -function in the last formula by the Euler's product formula, and define

$$\beta_{P,R}(n) := \sum_{\substack{D, v \geq 1 \\ Dv^2 = n^2 - 4 \\ p|v \Rightarrow p \leq P}} \left(\frac{1}{v} \prod_{p \leq P} \left(1 - \frac{\chi_D(p)}{p} \right)^{-1} \right) \prod_{p|d_B} \left\{ \begin{array}{l} 1 - \left(\frac{D}{p} \right), \quad p^2 \nmid D \\ 0, \quad p^2 \mid D \end{array} \right\}.$$

We have the local factor decomposition for this function [10], [9]:

Lemma 8.2 *Let $P \geq d_B$, then*

$$\beta_{P,R}(n) = \prod_{p \leq P} \beta_{(p,R)}(n),$$

where

$$\beta_{(p,R)}(n) := \sum_{b \geq 0} \frac{1}{p^b} \left(1 - \frac{1}{p} \chi_{(n^2-4)p^{-2b}}(p) \right)^{-1} \cdot \mathbb{I}_{p^b}(n).$$

Remark 8.3 *Here the definition of $\mathbb{I}_{p^b}(n)$ is*

$$\mathbb{I}_{2^b}(n) = \begin{cases} 1, & n^2 = 4 \pmod{2^{2b}}, (n^2 - 4)2^{-2b} \text{ is a discriminant} \\ 0, & \text{else} \end{cases};$$

$$\mathbb{I}_{p^b}(n) = \begin{cases} 1, & n^2 = 4 \pmod{p^{2b}}, p \neq 2 \\ 0, & \text{else} \end{cases};$$

$$\mathbb{I}_{p^b}(n) = \begin{cases} 1 - \left(\frac{(n^2 - 4)p^{-2b}}{p} \right), & n^2 = 4 \pmod{p^{2b}}, p^2 \nmid (n^2 - 4)p^{-2b} \\ 0, & \text{else} \end{cases}.$$

For the next result we use the uniform limit periodicity of $\beta_{(p,R)}$ and once again (see lemma 6.10) get the formula

$$\widehat{\beta_R}\left(\frac{a}{b}\right) = \prod_{p|b} \widehat{\beta_{(p,R)}}\left(\frac{a_p}{p^{\text{ord}_p b}}\right)$$

for the Fourier coefficients of β_R . It implies

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n) &= \sum_{b \geq 1} \sum_{\substack{1 \leq a \leq b \\ \gcd(a,b)=1}} \left| \widehat{\beta_R} \left(\frac{a}{b} \right) \right|^2 = \\ &= \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} \sum_{\substack{1 \leq a \leq p^c \\ a \not\equiv 0 \pmod{p}}} \left| \widehat{\beta_{(p,R)}} \left(\frac{a}{p^c} \right) \right|^2 \right), \end{aligned} \quad (13)$$

by using Parseval's equality.

Defining

$$\beta_{(p,R,b)}(n) := \left(1 - \frac{1}{p} \chi_{(n^2-4)p^{-2b}}(p) \right)^{-1} \cdot \mathbb{I}_{p^b}(n),$$

one obtain [10]

$$\widehat{\beta_{(p,R)}}(r) = \sum_{b \geq 0} \frac{1}{p^b} \widehat{\beta_{(p,R,b)}}(r).$$

For the Fourier coefficients in the right-hand side we have

Theorem 8.4 *For any prime $p \mid d_B$ the Fourier coefficients $\widehat{\beta_{(p,R,b)}}\left(\frac{a}{p^c}\right)$ are:*

$$\begin{aligned} c = 0, \quad b = 0, \quad \widehat{\beta_{(p,R,0)}}(0) &= \frac{(p-1)(p^2+2p+2)}{p^2(p+1)} \\ c = 0, \quad b \neq 0, \quad \widehat{\beta_{(p,R,b)}}(0) &= \frac{2(p^3-1)}{p^{2b+2}(p+1)} \\ c = 2b+2, \quad b \neq 0, \quad \widehat{\beta_{(p,R,b)}}\left(\frac{a}{p^c}\right) &= -\frac{2}{p^{2b+2}} \cos\left(\frac{4\pi a}{p^c}\right) \\ c = 2b+1, \quad b \neq 0, \quad \widehat{\beta_{(p,R,b)}}\left(\frac{a}{p^c}\right) &= -\frac{2}{p^{2b+2}(p+1)} \cos\left(\frac{4\pi a}{p^c}\right) - \\ &\quad - \frac{p^{1/2}\epsilon_p}{p^{2b}(p+1)} \left[e^{-4\pi i \frac{a}{p^c}} \left(\frac{-a}{p} \right) + e^{4\pi i \frac{a}{p^c}} \left(\frac{a}{p} \right) \right], \\ &\text{where } \epsilon_p = \begin{cases} 1, & p \equiv 1 \pmod{4} \\ i, & p \equiv 3 \pmod{4} \end{cases} \\ c \leq 2b, \quad b \neq 0, \quad \widehat{\beta_{(p,R,b)}}\left(\frac{a}{p^c}\right) &= \frac{2(p^3-1)}{p^{2b+2}(p+1)} \cos\left(\frac{4\pi a}{p^c}\right) \end{aligned}$$

$$c = 1, \quad b = 0, \quad \widehat{\beta_{(p,R,0)}}\left(\frac{a}{p}\right) = -\frac{2}{p^2(p+1)} \cos\left(\frac{4\pi a}{p}\right) - \frac{1}{p+1} \sum_{n \pmod{p}} \left(\frac{n^2-4}{p}\right) e^{-2\pi i n \frac{a}{p}}$$

$$c = 2, \quad b = 0, \quad \widehat{\beta_{(p,R,0)}}\left(\frac{a}{p^2}\right) = -\frac{2}{p^2} \cos\left(\frac{4\pi a}{p^2}\right).$$

Remark 8.5 The case $p \nmid d_B$ is done by M.Peter [10]. We will use these results later.

Corollary 8.6 For any prime $p \mid d_B$ the Fourier coefficients $\widehat{\beta_{(p,R)}}\left(\frac{a}{p^c}\right)$ are:

$$\widehat{\beta_{(p,R)}}(0) = 1$$

$$\widehat{\beta_{(p,R)}}\left(\frac{a}{p}\right) = \frac{-1}{p+1} \sum_{n \pmod{p}} \left(\frac{n^2-4}{p}\right) e^{-2\pi i n \frac{a}{p}}$$

$$\widehat{\beta_{(p,R)}}\left(\frac{a}{p^2}\right) = \frac{-2}{p(p+1)} \cos\left(\frac{4\pi a}{p^2}\right)$$

$$\widehat{\beta_{(p,R)}}\left(\frac{a}{p^c}\right) = \frac{-2}{(p+1)p^{\frac{3c-4}{2}}} \cos\left(\frac{4\pi a}{p^c}\right), \text{ for } 2 < c \text{ even}$$

$$\widehat{\beta_{(p,R)}}\left(\frac{a}{p^c}\right) = \frac{-\epsilon_p}{(p+1)p^{\frac{3c-4}{2}}} \left(\frac{a}{p}\right) \left[e^{-4\pi i \frac{a}{p^c}} \left(\frac{-1}{p}\right) + e^{4\pi i \frac{a}{p^c}} \right], \text{ for } 2 < c \text{ odd}$$

Now we able to calculate

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n).$$

Let us define the function

$$A_R(p^c) := \sum_{\substack{1 \leq a \leq p^c \\ p \nmid a}} \left| \widehat{\beta_{(p,R)}}\left(\frac{a}{p^c}\right) \right|^2.$$

Using (13) we find that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n) = \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} A_R(p^c) \right).$$

The values of the $A_R(p^c)$ for $p \nmid d_B$ were calculate by M.Peter [10]. They are:

$$p \neq 2, p \nmid d_B; \quad A_R(p) = \frac{p^2 - 2p - 1}{(p^2 - 1)^2}, \quad A_R(p^c) = \frac{2(p - 1)}{(p^2 - 1)^2 p^{2c-3}}$$

$$p = 2; \quad A_R(2) = \frac{1}{9}, \quad A_R(4) = \frac{1}{18},$$

$$A_R(8) = 0, \quad A_R(16) = \frac{1}{9 \cdot 16},$$

$$A_R(32) = 0, \quad A_R(2^c) = \frac{1}{9 \cdot 2^{2c-5}}, c \geq 6.$$

We complete his results by adding the case $p \mid d_B$. By simple calculations we have

Theorem 8.7 *Let $p \mid d_B$, then*

$$A_R(p) = \frac{p^2 - 2p - 1}{(p + 1)^2}; \quad A_R(p^2) = \frac{2(p - 1)}{p(p + 1)^2}; \quad A_R(p^c) = \frac{2(p - 1)}{(p + 1)^2 p^{2c-3}}.$$

Gathering the results we conclude

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n) &= \prod_{p \text{ - prime}} \left(1 + \sum_{c \geq 1} A_R(p^c) \right) = \\ &\left(1 + \frac{1}{9} + \frac{1}{18} + \frac{1}{9 \cdot 16} + \sum_{c \geq 6} \frac{1}{9 \cdot 2^{2c-5}} \right) \times \\ &\times \prod_{p \mid d_B} \left(1 + \frac{p^2 - 2p - 1}{(p + 1)^2} + \frac{2(p - 1)}{p(p + 1)^2} + \sum_{c > 2} \frac{2(p - 1)}{(p + 1)^2 p^{2c-3}} \right) \times \\ &\times \prod_{\substack{p \neq 2 \\ p \nmid d_B}} \left(1 + \frac{p^2 - 2p - 1}{(p^2 - 1)^2} + \sum_{c \geq 2} \frac{2(p - 1)}{(p^2 - 1)^2 p^{2c-3}} \right) \\ &= \frac{1015}{864} \prod_{p \mid d_B} \frac{2p(p^2 + p + 1)}{(p + 1)^3} \prod_{\substack{p \neq 2 \\ p \nmid d_B}} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p + 1)}. \end{aligned}$$

And finally we have

$$\begin{aligned}
\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{2 \leq n \leq N} \beta_R^2(n) &= \frac{1015}{864} \prod_{p|d_B} \frac{2p(p^2 + p + 1)}{(p + 1)^3} \prod_{\substack{p \neq 2 \\ p \nmid d_B}} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p + 1)} \\
&= \frac{1015}{864} \prod_{p|d_B} \frac{2p(p^2 + p + 1)}{(p + 1)^3} \cdot \frac{(p^2 - 1)^2(p + 1)}{p^2(p^3 + p^2 - p - 3)} \times \\
&\quad \times \prod_{p \neq 2} \frac{p^2(p^3 + p^2 - p - 3)}{(p^2 - 1)^2(p + 1)} \\
&= C_1 \cdot \prod_{p|d_B} \frac{2(p^3 - 1)(p - 1)}{p(p^3 + p^2 - p - 3)}, \quad \text{where } C_1 = 1.328...
\end{aligned}$$

This calculation concludes the proof.

References

- [1] Cohen.H.:A Course in Computational Algebraic Number Theory, Springer-Verlag, Berlin, 1993.
- [2] Cohn.H.:Advanced Number Theory, Dover Publications Inc.,1980.
- [3] Bogomolny.E., Leyvraz.F., Schmit.C.:Distribution of eigenvalues for the modular group. Comm. Math. Phys. 176 (1996), no. 3, 577-617.
- [4] De Bièvre.S: Quantum chaos: a brief first visit, Second Summer School in Analysis and Mathematical Physics (Cuernavaca, 2000), 161–218, Contemp. Math., 289, Amer. Math. Soc., Providence, RI, 2001.
- [5] Dedekind.K.:Über die Anzahl der Ideal-Klassen in den verschiedenen Ordnungen eines endlichen Körpers, in Gesammelte matematische Werke, Vol I, Chelsea Publishing Co., New York, 1968, 105-158.
- [6] Hejhal.D.A.: The Selberg Trace Formula for $PSL(2, \mathbb{R})$, vol. 1, 2, Lecture Notes in Math. Springer-Verlag, Berlin, 1976, 1983.
- [7] Ireland,K., Rosen,M.: A classical introduction to modern number theory, 2nd ed., Springer-Verlag, Berlin-Heidelberg-New York, 1990.
- [8] Landau,E.: Vorlesungen über Zahlentheorie, vols. 1-3, S.Hirzel, Leipzig,1927.
- [9] Lukianov.V.:Ph.D.Thesis (in preparation).
- [10] Peter.M.: The correlation between multiplicities of closed geodesics on the modular surface. Comm.Math.Phys. 225(2002), no.1, 171-189.
- [11] Peter.M: Momente der Klassenzahlen binärerquadratischer Formen mit ganzalgebraischen Koeffizienten. Acta Arith. 70, 43-77 (1995).
- [12] Peter.M: The limit distribution of a number theoretic functions arising from a problem in statistical mechanics. J. Number Th. 90, 265-280 (2001).
- [13] Miyake.T.:Modular forms, Springer-Verlag, New York, 1989.
- [14] Rudnick.Z.: A central limit theorem for the spectrum of the modular domain. Park City lectures 2002,to appear in Annales Henri Poincare.
- [15] Sarnak.P.: Class number of indefinite binary quadratic forms. J. Number Th.15, 229-247 (1982).
- [16] Strömbergsson.A.: Some remarks on a spectral correspondence for Maass waveforms. Internat. Math. Res. Notices 2001, no.10, 505-517.
- [17] Weil.A.:Basic Number Theory, 3rd ed., Springer-Verlag, New York, 1974.